
SHOUT
SHARE
IT

UNIVERSITY OF ILLINOIS
LIBRARY

Class
580.5

Book
F B

Volume
1
cop. 2

Ja 09-20M

CENTRAL CIRCULATION BOOKSTACKS
The person charging this material is responsible for its return to the library from which it was borrowed on or before the **Latest Date** stamped below.

Theft, mutilation, and underlining of books are reasons for disciplinary action and may result in dismissal from the University.

TO RENEW CALL TELEPHONE CENTER, 333-8400

UNIVERSITY OF ILLINOIS LIBRARY AT URBANA-CHAMPAIGN

APR 22 1992
APR 21 1992

When renewing by phone, write new due date below previous due date.

79521 L162

PUBLICATIONS

OF THE

FIELD COLUMBIAN MUSEUM

BOTANICAL SERIES

VOLUME I.

CHICAGO, U. S. A.
1895–1902.*

*For date of issue of each paper see Table of Contents.

FB
v. 1 cop. 2

1283
46494

24 '10 BINDING 1.09

TABLE OF CONTENTS.

PAGES.

Contribution to the Flora of Yucatan. (Aug., 1895.)
C. F. Millspaugh. (Plates I–IV.) 1– 56

Flora of West Virginia. (Jan., 1896.)
C. F. Millspaugh and L. W. Nuttall. (Plates V–VII.) 57–276

Contribution II to the Coastal and Plain Flora of Yucatan. (Dec., 1896.)
C. F. Millspaugh. (Plates VIII–XXI.) 277–339

Contribution III to the Coastal and Plain Flora of Yucatan. (Jan., 1898.)
C. F. Millspaugh 340–410

Higinbothamia. } (Aug., 1899.)
New Amaranthaceæ. }
E. B. Uline. (Plates XXII–XXIV.) 411–422

Plantæ Insulæ Ananasensis. (Aug., 1900.)
C. F. Millspaugh 423–440

Flora of the Island of St. Croix. (Nov., 1902.)
C. F. Millspaugh. (Map.) 441–550

150750

ERRATA.

A number of errors appear in the three Contributions to the Flora of Yucatan as published. These will be corrected in full in Vol. III of this series, which will be devoted to a descriptive Flora of Yucatan.

INDEX VOL. I.*

(ANTILLEAN SPECIES.)

For West Virginian Species see page 264.

Abelmoschus esculentus 507.
Abrus precatorius. 496.
Abutilon crispum 377.
graveolens 507.
incanum 377.
Indicum 29, 507.
lignosum 507.
sidoides 377.
umbellatum 507
Acacia Arabica 491
Catechu 491
cornigera 19
Farnesiana 19, 298, 363, 491.
filicina 298.
lutea 491.
spadicigera 19, 298, 363.
sphaerocephala 298.
tortuosa 491.
Vincentis 427.
Acalypha alopecuroides 27, 370.
mollis 302.
Portoricensis 502.
reptans 502.
setosa 27, 302, 371.
sp. 502.
unibracteata 302.
Yucatanense 371.
ACANTHACEÆ 46, 319, 391, 434, 529. 536.
Acanthophora muscoides 468.
Acanthospermum humile 533.
Achillea millefolium 394.
Achras Sapota 39, 313, 382, 517.
Achyranthes aspera 15, 294, 485.
aspera argentea 485.
aspera obtusifolia 485.
Acisanthera quadrata 431.
Acrocomia Mexicana 354.
Acrodiclidium salicifolium 489.
Acrostichum aureum 470.
calomelanos 469.
Adansonia digitata 510.
Adelia Oaxacana 401.
Ricinella 502.
Adenanthera pavonina 493.
Adiantum foliosum 469.
Adiantum fragile 469.
tenerum 348, 469.
tricholepis 8, 286, 348.
villosum. 469.
Adicea microphylla 483.
microphylla succulenta 483.
microphylla trianthemoides 483.
nummulariæfolia 483.
Æchmea bracteata 290, 355.
laxiflora 12, 290.
Ælidiella triumfettæ 466.
Ægiphila Martinicensis 523.
Æschynomene Americana 22, 298, 494.
Americana **depila** 363, 494,
fascicularis 298, 363.
tenuis 427.
Agardhiella tenera 468.
Agaricus **Yucatanensis** 285.
Agati grandiflora 494.
Agave Americana 480.
rigida elongata 292.
Sisalana 292.
sobolifera 480.
Ageratum conyzoides 323, 394, 533.
corymbosum 50, 323.
intermedium 51.
Agrostis Domingensis 474.
Indica 474.
Virginica 474.
AIZOACEÆ 487.
Albizzia Lebbek 491.
Albugo tragopogonis 465.
ALGÆ 347.
Aleurites Moluccana 503.
Alicastrum Brownei 14, 293, 358.
ALISMACEÆ 470.
Alisma cordifolia 470.
Allionia violacea 296, 361.
Allium fistulosum 479.
Allophyllus Cominia 402.
Aloe vera 292.
vulgaris 479.
Alomia ageratoides 51, 394.
Alpinia nutans 480.
Althernanthera Achyrantha 485.
Achyrantha 421.
asterotricha 419.
Culebrasensis 420.
Lorentzii 421.
Morongii 418.
obovata 360.
paronychioides 485.
polygonoides 485.
repens 360.
straminea 16, 295, 360.
Alvaradoa amorphoides 33, 301, 370.
Alysicarpus vaginalis 494.
AMARANTACEÆ 15, 294, 360, 418, 427, 485.
Amaranthus chlorostachys 15.
hybridus 15, 360.
paniculatus 486.
polygonoides 295, 486.
spinosus 295, 486.
tristis 486.
AMARYLLIDACEÆ 12, 292, 357, 426, 479.
Amaryllis equestris 12, 292.
tubispatha 479.
Ambrosia artemisiæfolia Trinitensis 533.
crithmifolia 52, 323.
hispida 52, 323.
Amellus aspera 52, 323.
niveus 52. 323.
niveus 395.
Amerimnon Brownei 363.
Campechianum 22.
Ammannia latifolia 513.
Amoreuxia palmatifida 380
Amyris sylvatica 500.
Anacanthus acicularis 529.
Jamaicensis 529.
spinosus 529.
ANACARDIACEÆ 32, 375, 504.

*New species in **black face** type.

Anacardium occidentale 375, 505.
Anamomis punctata 514.
Ananas sativus 12, 478.
Andropogon Halepensis 287.
hirtiflorus 350.
insularis 473.
Nardus 287.
saccharoides 471.
Schoenanthus 471.
semiberbis 350.
sorghum Drummondii 350.
Wrightii 471.
Aneimia adiantifolia 9, 287, 348.
bipinnata 348.
Angelonia angustifolia 389.
Anguria glomerata 532.
trilobata 532.
Anoda acerifolia 29.
hastata 29, 307, 377.
parviflora 377.
triangularis 377.
ANONACEÆ 17, 296, 361, 488.
Anona Cherimolia 296.
insularis 17.
laurifolia 488.
muricata 17, 361, 488.
palustris 361, 488.
reticulata 488.
squamosa 17, 361, 488.
Anthemis repens 53.
Anthephora elegans 287, 350, 473.
villosa 471.
Antheropogon triæna 10.
Antherylium Rohrii 513.
Anthurium cordifolium 477.
Huegelii 477.
Antigonon cordatum 484.
Antigonum leptopus 15, 359.
Antirrhœa lucida 530.
Aphelandra pectinata 46, 319.
APOCYNACEÆ 39, 313, 382, 432, 518.
AQUIFOLIACEÆ 429.
ARACEÆ 11, 355, 477.
Arachis hypogæa 363, 495.
ARALIACEÆ 432.
Ardisia coriacea 517.
Argemone Mexicana 18, 489.
Mexicana ochroleuca 297, 362.
Argithamnia candicans 502.
tinctoria 302.
Aristida Americana 350, 475.
dispersa 474.
nigrescens 350.
stricta 474.
tenuis 354.
ARISTOLOCHIACEÆ 15, 294, 359, 484.
Aristolochia anguicida 484.
brevipes 294, 359.
Aristolochia *fœtens* 15, 294.
grandiflora 15, 294.
maxima 359.
trilobata 484.
Artemisia Mexicana 323.
Artocarpus incisa 358, 482.
Arum Italicum 355.
maculatum 478.
Arundo donax 350.
ASCLEPIADACEÆ 36, 313, 384, 433, 519, 537.
Asclepias Curassavica 36, 313, 384, 519.
longicornu 314.
pratensis 38.
Asemnantha pubescens 48.
Asparagus officinalis 292.
Aspergillus flavus 285.
fuliginosus 285.
Aspidium exaltatum 468, 469.
invisum 468.
molle 468.
patens 468.
trifoliatum 9, 348.
Asplenium dentatum 349.
pumilum 8. 287, 349.
Asterina colubrinæ 465.
Yucatanensis 285.
Aster Novi-Belgii 323.
Atriplex cristata 15, 359, 485.
Ateleia microcarpa 21.
pterocarpa 21.
Attalea Cohune 355.
Avicennia nitida 42, 316, 386, 523.
officinalis 42, 316, 386.
tomentosa 523.
tomentosa 42, 316, 386.
Ayenia magna 31, 310.
ovata 310.
pusilla 310, 379, 510.
Yucatanense 379.

Baccharis dioica 533.
BALSAMINACEÆ 506.
Bambusa vulgaris 475.
Barbula agraria 7.
Bastardia viscosa 507.
BATIDACEÆ 486.
Batis maritima 486.
Bauhinia Cavanillei 364.
divaricata 20, 298.
Lamarckiana 20.
latifolia 20, 364.
porrecta 364.
porrecta 20, 298.
splendens 364.
tomentosa 492.
Beloperone violacea 320.
Belotia grewiæfolia 430.
Benthamantha Caribæa 494.
Bidens Alansensis 54.
bipinnata 533.
Bidens leucantha 323, 394, 533.
nivea 52.
pilosa 394.
tereticaulis 394.
BIGNONIACEÆ 45, 390, 319, 434, 528.
Bignonia æquinoctialis 390, 528.
diversifolia 390.
mollis 390.
Stans 45.
Unguis 528.
Unguis-cati 390.
BIXACEÆ 32, 311, 380, 511.
Bixa Orellana 32, 511.
Blainvillea dichotoma 52.
Blechum blechoides 320.
Blechum 529.
Brownei 46, 320.
laxiflorum 320.
occidentale 469.
Boerhaavia erecta 17, 295, 361, 486.
hirsuta 17, 296.
paniculata 487.
scandens 361.
BOMBACACEÆ 30, 309, 379, 510.
Bombax Ceiba 30.
Ceiba 30.
pentandrum 30.
BORRAGINACEÆ 40, 315, 385, 433, 522.
Borreria thymifolia 49.
parviflora 530.
stricta 530.
verticillata 530.
Borrichia arborescens 53, 533.
argentea 533.
Bouchea Ehrenbergii 523.
Ehrenbergii 41.
prismatica 41, 391.
Bougainvillea spectabilis 487.
Boussingaultia baselloides 485.
Boutelona Americana 350.
bromoides 350.
disticha 350.
litiginosa 475.
triæna 10, 350.
trianthera 10.
Bradburya pubescens 23.
Schottii 364.
Virginiana 23, 364, 496.
Virginiana angustifolia 496.
Brassica juncea 489.
oleracea 297.
Bravaisia tubiflora 46, 391.
Brickellia diffusa 394.
BROMELIACEÆ 11, 290, 355, 478.
Bromelia bracteata 290.
Pinguin 11, 291, 355, 478.
sylvestris 291.
Bromus virgatus 352.
Brosimum Alicastrum 14.

Brunfelsia Americana 526.
Americana pubescens 526.
Brya Ebenus 427.
Bryophyllum calycinum 18, 298.
pinnatum 18, 298, 362, 490.
Bryothamnion Seaforthii 468.
Buceras Buceras 515.
Catappa 36, 312, 381, 516.
Buchnera Mexicana 45, 389.
Bucida Buceras 431.
Buceras 515.
Buettneria Carthaginensis 379.
Bumelia buxifolia 376.
cuneata 517.
glomerata 376.
microphylla 376.
obtusifolia buxifolia 376.
retusa 37.
Bunchosia glandulosa 369.
Lindeniana 24.
Swartziana 497.
Buphthalmium arborescens 53,
BURSERACEÆ 25, 302, 370, 428. 500.
Bursera graveolens 302.
gummifera 500.
gummifera 25.
Simaruba 25.
BUXACEÆ 504.
Buxus Vahlii 504.
Byrsonima coriacea 497.
crassifolia 24.
Cubensis 429.
Karwinskiana 24.
spicata 497.
verbascifolia 428.

Cabomba aquatica 17.
Cacara erosa 300.
CACTACEÆ 35, 311, 513.
Cactus Opuntia 35.
Opuntia 35.
Cæsalpinia Bonducella 21, 364.
coriaria 493.
Crista 493.
divergens 493.
exostemma 21.
melanosperma 493.
pulcherrima 21, 298, 364, 493.
Cajanus Cajan 496.
Cakile æqualis 18, 297, 489.
æqualis 362.
maritima 362.
maritima 18.
Caladium bicolor 355, 477.
Calea aspera 52, 323.
axillaris urticifolia 394.
Zacatechichi 394.
Calliandra gracilis 298, 364.
Portoricensis 19, 298, 364.
purpurea 491.
Callicarpa acuminata 42, 316.
Callicarpa reticulata 523.
Callisia repens 13, 356, 479.
Calophyllum Calaba 511.
Calopogonium cæruleum 23.
Calotropis procera 519.
Calycorectes Mexicanus 312.
Calyptranthes punctata 431.
Calyptrion Aubletii 32.
Hybanthus 32.
Calyptronoma Swartzii 426.
Cameraria retusa 432.
CAMPANULACEÆ 50, 322, 533.
Canavalia Cubensis 428.
ensiformis 365.
obtusifolia 496.
CANELLACEÆ 511.
Canella alba 511.
CANNACEÆ 13, 358. 481.
Canna edulis 358, 481.
Indica 13, 481.
Lamberti 481.
lutea 481.
CAPPARIDACEÆ 18, 297, 362, 427, 489.
Capparis amygdalina 18, 297.
cynophallophora 362, 490.
frondosa 490.
Grisebachii 490.
Jamaicensis 490.
Capraria biflora 45, 319, 389, 527.
biflora pilosa 45, 527.
saxifragæfolia 390.
CAPRIFOLIACEÆ 49, 321, 393, 532.
Capsicum annuum 388, 526.
baccatum 43, 318, 526.
dulce 526.
frutescens 43, 318, 388, 526.
Cardiospermum Corindum
molle 402.
Halicacabum 505.
microcarpum 505.
molle 306.
CARICACEÆ 34, 381. 512.
Carica Papaya 34, 381, 512.
Carolinea alba 30, 309.
fastuosa 30, 379.
macrocarpa 30.
CARYOPHYLLACEÆ 296, 488.
Casearia hirta 511.
parviflora microcarpa 511.
ramiflora 512.
sylvestris 512.
Cassia alata 20, 492.
bacillaris 20.
bicapsularis 20, 492.
biflora 20, 365.
biflora angustisiliqua 492.
chamæcrista 298.
emarginata 20, 298, 365.
fistula 492.
flavicoma 365.
grandis 492.
Cassia hirsuta 365.
leptocarpa hirsuta 365.
Liebmanni 365.
obtusifolia 492.
occidentalis 20, 298, 365, 492.
polyphylla 366, 492.
procumbens 366.
racemosa 366.
ramosa 493.
reticulata 20.
sericea 299, 366.
Tora 493.
tristicula 366.
villosa 366.
Cassimiroa **tetrameria** 401.
Cassytha Americana 489.
Americana 18.
filiformis 18.
Castalia ampla 17, 296, 361, 488.
Castela erecta 500.
Castilleja arvensis 319.
communis 319.
Castilloa elastica 358.
CASUARINACEÆ 14, 358, 482.
Casuarina equisetifolia 14, 358, 482.
Catesbæa melanocarpa 530.
CAULERPACEÆ 467.
Caulerpa clanifera 467.
plumaris 467.
Cayaponia alata 50, 322, 393.
graciliflora 532.
grandiflora 50, 393.
Cecropia obtusa 358.
peltata 482.
Cedrela odorata 26, 302, 370.
Glaziovii 26.
Cedronella Mexicana 387.
Ceiba casearia 308, 379.
pentandra 30.
Schottii 308, 379.
CELASTRACEÆ 32, 375, 401, 505.
Celosia argentea 485.
nitida 485.
nitida 15.
paniculata 15, 295.
virgata 295.
Celtis aculeata 399, 482.
aculeata serrata 482.
trinervis 482.
Cenchrus brevisetus 10.
echinatus 351, 474.
pallidus 9, 351.
viridis 474.
Centrosema pubescens 23.
Virginianum 23.
Cephalandra Indica 532.
Cephalanthus occidentalis salicifolius 435.
Cereus flagelliformis 35.
floccosus 513.
grandiflorus 35, 513.
nycticalus 513.

Cereus pectem aboriginum 311.
triangularis 513.
Cestrum diurnum 44, 318, 388, 526.
laurifolium 526.
nocturnum 526.
CHÆTANGIACEÆ 468.
Chætochloa caudata 473.
glauca 473.
setosa 473.
Chætolepis Cubensis 431.
saturæioides 431.
CHÆTOPHORACEÆ 467.
Chalcas exotica 25.
Chamæcrista glandulosa 493.
glandulosa ramosa 493.
nictitans 493.
Chamædorea gracilis 355.
Chamæraphis glauca 473.
Chamissoa paniculata 360.
Chaptalia nutans 533.
CHARACEÆ 286, 347.
Chara gymnopus 347.
gymnopus inconstans 286.
Cheilanthes leucopoda 8.
microphylla 287, 349, 469.
CHENOPODIACEÆ 15, 359, 485.
Chenopodium album 359.
ambrosioides 485.
murale 485.
Chiococca racemosa 48, 321, 392, 530.
Chione glabra 530.
Chloris barbata 351.
ciliata 351, 474.
elegans 351.
eleusinoides 474.
petræa 10.
radiata 475.
Chlorophora tinctoria 358, 482.
CHLOROPHYCEÆ 286.
Chrysobalanus Icaco 19, 490.
Chrysodium vulgare 470.
Chrysophyllum Cainito 313, 382, 517.
glabrum 517.
microphyllum 518.
olivæforme 518.
Chytraculia Chytraculia 514.
pallens 514.
punctata 431.
Cinnamomum Zeylanicum 489.
Cipura plicata 480.
Cissampelos Pareira 297, 362, 427, 488.
Pareira microcarpa 488.
Cissus acida 506.
sicyoides 506.
trifoliata 506.
Citharexylum cinereum 524.
quadrangulare 386, 524.
Citrus acida 301.
Citrus Aurantium 25, 301, 499.
buxifolia 499.
decumana 499.
Limetta 25, 301, 499.
Limonum 25, 499.
medica 499.
vulgaris 25, 301, 499.
Cladium germanicum 290, *mariscus* 290.
Clematis Caracasana 17.
Caripensis 17.
dioica 17, 297.
flammulastrum 17.
Cleome pentaphylla 489.
polygama 297.
procumbens 427.
pungens 490.
spinosa 297, 362, 490.
Clerodendron aculeatum 316, 386, 524.
Colebrookianum 386.
fragrans 316.
ligustrinum 42.
siphonanthus 524.
Clidemia petiolaris 36.
spicata 432, 516.
Clinopodium Brownei 43, 317.
Clitoria cajanifolia 428.
glomerata 428.
Mexicana 366.
ternatea 496.
Virginiana 23.
Clusia alba? 511.
rosea 511.
CLYPEOSPHÆRIACEÆ 465.
Cnicus Mexicanus 323.
Coccoloba Barbadensis 484.
coronata 294, 484.
Cozumelensis 15.
diversifolia 484.
Klotzschiana 537.
laurifolia 484.
leoganensis 484.
microstachya ovalifolia 484.
nivea 484.
obtusifolia 484.
punctata parviflora 484.
retusa 427.
uvifera 15, 294, 484.
Cochlospermum hibiscoides 31, 311.
Cocos nucifera 11, 290, 355.
CODIACEÆ 467.
Codiæum variegatum 371, 502.
Codium tomentosum 467.
Coffea Arabica 392.
Liberica 530.
Coix lachryma-Jobi 9, 351, 470.
Coleosporium Ipomœæ 465.
Coleus Amboinicus 525.
Colubrina ferruginosa 376, 506.
reclinata 506.
COMBRETACEÆ 36, 312, 381, 431, 515.
COMMELINACEÆ 13, 292, 356, 478.
Commelina erecta 13.
nudiflora 356, 478.
pallida 356.
Virginica 13, 478.
Comocladia ilicifolia 504.
COMPOSITÆ 50, 323, 394, 435, 533.
Condalea ferrea 506.
Conobea pusilla 389.
Conocarpus erectus 36, 516.
erectus arboreus 312.
erectus procumbens 516.
sericea 381.
Conradia pumila 434.
Conostegia Xalapensis 36.
CONVOLVULACEÆ 39, 314, 384, 433, 519.
Convolvulus Grayi 314.
Jamaicensis 40, 519.
Jamaicensis 39.
nodiflorus 519.
pentanthos 39.
Conyza lobata 54.
odorata 52.
purpurascens 52.
Corallocarpus emetocathartic us 322.
glomeratus 322.
Millspaughi 322, 393.
Corchorus acutangulus 507.
hirsutus 507.
hirtus 507.
pilolobus 307.
siliquosus 34, 376, 507.
Cordia alba 522.
Collococca 522.
cylindristachya 315, 522.
dodecandra 40, 315, 385.
gerascanthoides 40, 385.
gerascanthus 40.
globosa 40, 315, 385, 522.
Martinicensis 522.
nitida 522.
Rickseckeri 522.
Sebestena 40, 315, 386, 522.
speciosa 40.
sulcata 522.
ulmifolia 522.
ulmifolia lineata 522.
Cornutia pyramidata 316.
Corynostylis Hybanthus 32.
Cosmos caudatus 394, 533.
Cotyledon pinnatum 18.
Couroupita Guianensis 514.
Coussarea speciosa 48.
Coutarea octomera 47.
Cracca bicolor 366.
Caribæa 22.
cinerea 22, 299.
Greennanii 299, 366.

Craca villosa cinerea 366, 494.
CRASSULACEÆ 18, 298, 362, 490.
Cratæva gynandra 297, 362.
Tapia 18.
Tapia 297.
Crescentia cucurbitina 528.
Cujete 45, 319, 390, 528.
Crinum erubescens 426, 479.
giganteum 479.
longiflorum 480.
Crossandra infundibuliformis 529.
Crossopetalum pallens 32.
Crotalaria incana 299, 494.
juncea 494.
latifolia 494.
pumila 21, 366.
retusa 494.
verrucosa 494.
Croton albidus 303.
arboreus 303.
astroites 501.
ciliato-glandulosus 371.
Cortesianus 303, 371.
betulinus 501.
bispinosus 429.
cerinus 429.
crasspedotrichus 429.
discolor 429, 501.
flavens 303, 371, 501.
flavens rigidus 501.
fragilis 303.
humilis 303, 371.
lobatus 303, 371, 429, 501.
maritimus 26, 371.
ovalifolius 501.
reptans 429.
rhamniæfolius salviæfolius 303.
Zalapensis 26.
CRUCIFERÆ 18, 297, 362, 489.
Crusea calocephala 321, 392.
Cryphæa filiformis 286.
Cucumis Anguria 393, 532.
CUCURBITACEÆ 50, 322, 393, 435, 532.
Cucurbita ficifolia 532.
Lagenaria 532.
Pepo 532.
Pepo Melopepo 532.
Cuphea Pseudomelanium 431.
Trinitatis 311.
Curcuma longa 480.
Cuscuta Americana 40, 314, 519.
Cyathus Pœphegii 467.
CYCADACEÆ 9.
Cyclanthera ribiflora 322.
Cymodocea nanatorium 470.
Cynanchum parviflorum 38.
racemosum 38.
Cynodon Dactylon 287, 351, 474.
Cynosurus Ægyptiacus 351, 475.
Indicus 475.
Cynosurus virgatus 475.
CYPERACEÆ 10, 290, 354, 475, 537.
Cyperus aristatus 10.
articulatus 476.
Brizæus 476.
Brunneus 10, 476.
canus 290.
capillaris 476.
confertus 476.
Eggersii 476.
elegans 476.
filiformis 476.
flavomariscus 476.
flavus 476, 537.
interstinctus 425.
lævigatus 476.
ligularis 11, 476.
Michauxianus 476.
ochraceus 290, 354, 476.
Ottonis 11.
rotundus 354, 477.
squarrosus 354.
tenuis 537.
thyrsiflorus 11.
viscosus 476.
Cystopus tragopogonis 465.

Dactyloctenium Ægypticum 351, 475.
Dalbergia Amerimnum 363.
Campechiana 22.
Ecastaphyllum 495.
Dalea diffusa 366.
Domingensis 21, 299.
Dalechampia scandens 304, 371, 502.
Dasylirion pliabile 12.
Datura fastuosa 526.
Metel 318, 526.
Stramonium 44, 388, 526.
Tatula 527.
Desmanthus depressus 20, 367.
virgatus 492.
virgatus strictus 492.
Desmodium affine 22.
scorpiurus 299.
tortuosum 299.
Dianthera comata 47.
pectoralis 529.
sessilis 47.
sessilis 320.
Diapedium assurgens 46, 320, 391, 529.
Dicliptera assurgens 46, 320, 391.
Didymodon æneus 7.
DICTYOTACEÆ 467.
Dictyota ciliata 467.
dentata 467.
Digitaria marginata 353, 473.
setigera 473.
Dilophus alternans 468.
Diodia teres 321.
Dioon spinulosum 9.
DIOSCOREACEÆ 13, 359, 413, 426, 480.
Dioscorea alata 357, 480.
altissima 480.
calyculata 357.
densiflora 13.
floribunda 357.
macrostachya 357.
Matagalpensis 417.
polygonoides 357.
sativa 357.
spiculiflora 13.
trifida 426.
Yucatanensis 416.
Diospyros Ebenum 382.
laurifolia 432.
Dipholis salicifolia 382, 518.
Diphysa Carthaginensis 367.
robinioides 367.
Diplachne fascicularis 475.
Diplochita serrulata 516.
Distichlis spicata 351.
Distictis lactiflora 528.
Distreptus spicatus 534.
Ditaxis fasciculata 501.
Dodonæa viscosa 506.
Dolicholus minimus 496.
minimus luteus 496.
reticulatus 496.
Dolichos Lablab 497.
Lablab albiflorus 497.
minima 23.
pruriens 23.
Dorstenia Contrajerva Houstoniana 293, 358.
DOTHIDACEÆ 465.
Drejerella mirabiliodes 529.
Drepanocarpus lunatus 495.
Drymaria cordata 488.
Dryopteris patens 287.
Drypetes glauca 501.
lævigata 501.
Duranta Plumieri 386.
Plumieri 42, 316.
repens 42, 316, 524.
spinosa 386.
Xalapensis 316.
Dysodia cancellata 323, 395.
Dysodium divaricatum 53.

EBENACEÆ 382, 432.
Echeandia terniflora 292.
Echinodorus cordifolius 470.
rostratus 470.
Echites adglutinata 518.
barbata 518.
biflora 382.
microcalyx 313.
microcalyx 382.
neriandra 518.
paludosa 433.

Echites suberecta 518.
subsagittata 382.
torulosa 383.
umbellata 383, 433.
Eclipta alba 395, 534.
erecta 323.
Egletes viscosa 52.
Ehretia Bourreria 522.
tinifolia 40, 315, 386.
Eichhornia azurea 479.
Elæis occidentalis Sw 426.
Elæodendron xylocarpum 505.
Eleocharis capitata 290.
capitata 477.
cellulosus 477.
interstincta 477.
nodulosa 477.
Elephantopus mollis 534.
Eleusine Ægyptica 10.
Ægyptica 475.
Indica 10, 288, 351, 475.
Eleutheranthera divaricata 53, 324.
Elvira biflora 323.
Martyni 323.
Elytraria frondosa 320.
tridentata 46, 320.
Emilia sagittata 534.
sonchifolia 534.
Enslenia racemosa 38.
Epidendrum bifidum 481.
ciliare 481.
cochleatum 481.
pallidiflorum 426.
Eragrostis ciliaris 10, 288, 351, 475.
elongata 351.
Mexicanus 288.
minor 475.
pilosa 475.
plumosa 351.
Eranthemum nervosum 536.
Erechthites hieracifolia 534.
hieracifolia cacalioides 534.
Erigeron atriplicifolium 534.
Ayapana 534.
Canadensis 52, 323, 395.
spathulatus 534.
Erinus procumbens 44, 319.
Eriochloa punctata 471.
Eriodendron anfractuosum 510.
anfractuosum 30.
Eriosema crinitum 428.
Erithalis fruticosa 48, 530.
fruticosa odorifera 530.
Ernodea littoralis 49, 321, 392, 530.
Erythrina carnea 23.
Corallodendron 367, 496.
coralloides 299, 367.
horrida 496.
Piscipula 22.
Erythroxylaceæ 428, 497.
Erythroxylum alaternifolium 428.
brevipes 497.
ovatum 497.
Esenbeckia pentaphylla 24, 401.
Ethulia ageratoides 51.
Eucharis grandiflora 480.
Eugenia acetosans 514.
axillaris 514.
brevipes 431.
buxifolia 514.
dichotoma 431.
floribunda 514.
glabrata 514.
lateriflora 514.
ligustrina 514.
monticola 515.
pallens 515.
Poiretii 515.
Portoricensis 515.
procera 515.
pseudopsidium 515.
sessiliflora 515.
uniflora 515.
virgultosa 515.
Eulejunea lepida 7.
Eupatorium albicaule 323.
conyzoides 51, 395.
drepanophyllum 324.
fœniculaceum 435, 534.
Houstonis 51.
macrophyllum 534.
odoratum 51, 534.
paniculatum 51.
populifolium 51, 324.
pycnocephalum 395.
quadrangulare 324.
repandum 534.
sinuatum 534.
Euphorbiaceæ 26, 302, 370, 429, 500.
Euphorbia adenoptera 28, 304, 371.
arenaria 371.
Armourii 28, 372.
articulata 503.
astroites 304.
astroites **heterappendiculata** 28.
buxifolia 28, 372, 503.
cyathophora 503.
dentata 372.
Gaumeri 372.
geniculata 503.
graminea **lancifolia** 372.
graminea **virgata** 372.
heterophylla 29, 305, 373.
heterophylla cyathophora 305, 373.
hypericifolia 27, 305, 503.
Euphorbia hypericifolia hyssopifolia 503.
lasiocarpa 373.
maculata 27.
Mayana 305, 373.
neriifolia 503.
ovalifolia 27.
petiolaris 29, 503.
pilosula 373.
pilulifera 17, 503.
pilulifera procumbens, 305, 373, 504.
Preslii 305.
prostrata 504.
pulcherrima 374.
rhytisperma 305, 374.
serpyllifolia 305, 374.
thymifolia 305, 504.
trichotoma 29.
Xbacensis 374.
Eustoma exaltatum 37.
Euxolus caudatus 486.
oleraceus 486.
Evolvulus alsinoides 314, 384.
linifolius 520.
mucronatus 520.
nummularius 39, 520.
sericeus 433.
Excœcaria glandulosa 305.
Exostemoma Caribæa 392, 531.
ellipticum 434.

Fagara microphylla 499.
Faramea odoratissima 435, 531.
Festuca fascicularis 475.
Fevillea cordifolia 435.
Ficus Bonplandiana 358.
crassinervia 483.
grandifolia? 14.
Jaliscana 293.
lævigata 483.
laurifolia 14.
longipes 293, 359.
pedunculata 483.
populinea 483.
rubiginosa? 14.
trigonata 483.
Filices 8, 286, 348.
Fimbristylis diphylla 477.
polymorpha 11.
spadicea 477.
Fischeria crispiflora 519.
Flacourtiaceæ 380, 431, 511.
Flacourtia inermis 431.
Flaveria linearis 54, 395.
longifolia 54.
repanda 295.
Fleurya æstuans 483.
Fœniculum vulgare 312.
Forchhammèria **trifoliata** 399.
Forestiera Jacquinii 518.
Forsteronia corymbosa 433.
Fourcroya Cubensis 480.

Fruillania ericoides 286.
FUCACEÆ 467.
FUNGI 285, 347.

Galactia angustifolia retusa 428.
filiformis 428.
filiformis 538.
multiflora 367.
regularis 495.
striata tomentosa 537.
tenuiflora 495.
Galaxaura fragilis 468.
Galega cinerea 22.
Galphimia glauca 498.
Gaudichaudia filipendula 369.
GENTIANACEÆ 37.
Gentiana exaltata 37.
Geonoma Swartzii 426.
Gerardia Domingensis 434.
GESNERIACEÆ 434.
Gilibertia cuneata 432.
Gliricidia maculata 22, 300.
Glœosporium affine 285.
Glycine labialis 496.
Gomphrena capituliflora 16.
decumbens 295, 360.
globosa 360, 485.
vermicularis 16.
Gonolobus barbatus 314.
GOODENIACEÆ 50, 394, 533.
Gossypium Barbadense 507.
herbaceum 377.
religiosum 377.
Gouania Domingensis 376, 506.
stipularis 33.
Gouinia latifolia 352.
polygama 352.
virgata 352.
GRAMINEÆ 9, 287, 350, 470.
Graptophyllum pictum 537.
Gratiola Monniera 44.
Gronovia scandens 35, 311, 381.
Guadua latifolia 352.
Guajacum officinale 498.
sanctum 428.
Guazuma polybotria 379.
tomentosa 31, 310.
ulmifolia 510.
Guettarda calyptrata 435.
elliptica 321.
parviflora 531.
scabra 48, 531.
Guignardia sp 465.
Guilandina Bonducella 21.
melanosperma 493.
Moringa 490.
GUTTIFERÆ 31, 380, 430, 511.
Gymnanthes lucida 374, 503.
Gymnogramma calomelanos 469.
sulphurea 469.

Gynoxis Berlandieri 396.
Gyrocarpus Americanus 381.

Hæmatoxylon Campechianum 21, 300, 493.
Hæmocharis Curtyana 430.
HÆMODORACEÆ 426.
Hamelia chrysantha 48.
lutea 531.
patens 48, 321, 392, 531.
Hechtia Schottii 12.
Hedwigia balsamifera 500.
Hedysarum adsendens 22.
Hedysotis Americana 47.
Helenium quadridentatum 324.
Heleocharis capitata 477.
cellulosa 477.
interstincta 425, 477.
nodulosa 477.
Helianthus annuus 395.
Helicteres Baruensis 310, 379.
Jamaicensis 510.
Heliotropium Currassavicum 315, 386, 433, 523.
fruticosum 386, 523.
gnaphalodes 41.
Indicum 523.
inundatum 41, 315, 386.
parviflorum 41, 315, 386, 523.
Peruvianum 523.
phyllostachyum 315.
reversiferum 433.
Hemionitis palmata 349, 469.
Hemitelia Hartii 8.
Henriettella parviflora 432.
Henrya costata 391.
costata 47, 320.
HEPATICÆ 7, 286.
Heptanthus ranunculoides 435.
Herpestis chamædryoides 527.
chamædryoides 44, 300.
Monniera 44.
Heteranthera limosa 292.
Heteropteris Beechyana 301, 369.
floribunda 24.
Lindeniana 24.
longifolia 24.
parviflora 498.
purpurea 498.
Yucatanensis 369,
Hibiscus Berlandierianus 307.
Braziliensis 508.
clypeatus 508.
mutabilis 378.
phœniceus 508.
Rosa-sinensis 307.
Sabdariffa 508.
tubiflorus 30, 307, 378.
vitifolius 508.
Higinbothamia synandra 415.
Hippeastrum equestre 12, 292, 480.

HIPPOCRATEACEÆ 33, 306, 375, 402, 430.
Hippocratea celastroides 306.
Grisebachii 402.
obcordata 375.
tenella 33.
verrucosa 402.
Hippomane Mancinella 503.
Hiræa Barclayana? 23.
Hirtella mollicoma 427.
Hisingera nitida 32.
Holmskioldia sanguinea 524.
Huertea Cubensis 430.
Hura crepitans 503.
polyandra 374.
Hybanthus **Yucatanensis** 404.
HYDROCHARITACEÆ 470.
Hydrocotyle prolifera 381.
Hydrolea glabra 40.
HYDROPHYLLACEÆ 40, 315, 521.
Hymenæa Courbaril 492.
Hymenocallis Caribæa 480.
lacera 292.
rotata 292.
HYMENOMYCETACEÆ 467.
Hypericum galioides Cubensis 430.
Hypnea musciformis 468.
Hypospila cordiana 465.
Hypoxis juncea 426.
scorzoneræfolia 426.
Hyptis capitata 42.
capitata 42, 525.
pectinata 42, 387.
pectinata 525.
suaveolens 387.
suaveolens 525.

Ibatia maritima 519.
Ichnanthus **lanceolatus** 352.
Ichythyomethia Piscipula 22, 300, 495.
Ilex montana 429.
Impatiens Balsaminea 506.
Indigofera Anil 21, 300, 494.
mucronata 21.
pascuorum 427.
tinctoria 494.
Inga laurina 491.
Ionidium brevicaule 311.
oppositifolium 311.
riparium 311.
strictum 511.
Ipomœa argentifolia 433.
Batatas 433, 520.
Batatas porphyrorhiza 520.
biloba 314.
bona-nox 39, 314, 384, 520.
carnea 520.
cissoides 384.
coccinea 39, 314, 384.
eustachyana 520.
fastigiata 39, 384, 520.

Ipomœa filiformis 520.
hederacea 384.
Jalapa 39.
Jamaicensis 39, 314.
mollicoma 520.
pentaphylla 385, 520.
Pes-capræ 39, 520.
Pes-capræ 314.
puncticulata 39, 385.
quinquepartita 520.
repanda 433.
sidæfolia 39.
sinuata 385, 520.
Steudelii 520.
triloba 39, 521.
triquetra 521.
Tuba 521.
tuberosa 521.
violacea 521.
Iresine celosioides 16, 295, 360.
elatior 485.
Hartmanii 422.
luzuliflora 16.
Iria polymorpha 11.
IRIDACEÆ 13, 426, 480.
Isocarpha oppositifolia 395.
Isotoma longiflora 50, 322, 533.
Ixophorus glaucus 473.
Ixora stricta 531.

Jacaratia Mexicana 35.
Jacquemontia abutiloides 314, 385.
pentantha 39, 314, 385, 521.
tamnifolia 385, 521.
violacea 39.
Jacquinia aristata 36.
armillaris 312, 382, 517.
Caracasana 36.
Jambosa Jambos 515.
Malaccensis 515.
Jasminum humile 518.
officinale 518.
pubescens 518.
quinqueflorum 518.
Jatropha aconitifolia 27.
Curcas 305, 374, 502.
gossypifolia 502.
gossypifolia elegans 503.
gossypifolia staphisagriæfolia 503.
multifida 429, 503.
urens 305, 374.
urens stimulosa 305, 374.
Juncellus lævigatus 476.
lævigatus albidus 476.
Jussieua angustifolia 516.
octovalvis 516.
suffructicosa 312, 517.
Justicia assurgens 46.
Carthaginensis 529.
pectoralis 529, 536.
Justicia periplocæfolia 536.
secunda 536.

Kallstrœmia maxima 498.
Karatas Plumieri 356.
Karwinskia Humboldtiana 376.
Kokera paniculata 360, 485.
Kyllinga brevifolia 425, 475.
filiformis 475.
monocephala 475, 476.
obtusata 476.
pumila 475.
pungens 476.
Krugiodendron ferreum 506.

LABIATÆ 42, 316, 387, 434, 525.
Lablab vulgaris albiflorus 497.
Lachnorrhiza piloselloides 435
Lactuca Intybacea 534.
Læstadia Rickseckeri 465.
Lagascea mollis 324.
Lagenaria Lagenaria 534.
Lagenaria viscosa 532.
Lagerstrœmia Indica 513.
Lagetta lintearia 431.
Laguncularia racemosa 431, 516.
Lantana aculeata 524.
Camara 41, 316, 386, 524.
involucrata 41, 316, 434, 524.
odorata 41.
odorata 316.
polyacantha 524.
reticulata 524.
Laplacea Curtyana 430.
Lappago aliena 471.
LAURACEÆ 17, 297, 362, 427, 489.
Laurencia obtusa 468.
Lawsonia alba 35.
inermis 514.
LECYTHIDACEÆ 514.
LEGUMINOSÆ 19, 298, 363, 427, 491, 537.
LEMNACEÆ 290, 355, 478.
Lemna minor 478.
LENNOACEÆ 382.
Lennoa cærulea 382.
Lentinus villosus 347.
Leonotis nepetæfolia 525.
Leonurus glaucescens 387.
Sibiricus 525.
Lepidium apetalum 297.
Virginicum 489.
Leptilon Canadense 535.
Leptochloa fascicularis 475.
mucronata 288, 352, 475.
virgata 475.
virgata gracilis 475.
virgata multiflora 475.
Leucæna glauca 19, 300, 491.
Leucas Martinicensis 525.
Leucobryum **incurvifolium** 347.
LICHENES 347.
LILIACEÆ 12, 292, 357, 426, 479.
Lippia Berlandieri 316.
dulcis 316.
geminata 317.
graveolens 317.
nodiflora 42, 317, 434, 524.
origanoides 42.
stœchadifolia 42, 317.
umbellata 387.
Lithophila muscoides 485.
vermiculata 427, 485.
LOASACEÆ 35, 311, 381.
Lobelia longiflora 50.
Plumieri 50.
LOGANIACEÆ 432.
Lonchocarpus violaceus 22.
Lonicera Japonica 532.
macrantha 393.
LORANTHACEÆ 294, 483.
Loranthus calyculatus 294.
Lourea vespertilionis 495.
Lucuma mammosa 37.
multiflora 518.
multifida 313.
Lueha speciosa 376.
Luffa Ægyptica 393.
cylindrica 50, 532.
cylindrica 393.
Lycopersicum cerasiforme 527.
esculentum 527.
Humboldtii 44.
LYCOPODINÆ 287, 349, 468.
Lysiloma latisiliqua 300.
Sabicu 300.
LYTHRACEÆ 35, 311, 431, 513.

Maba Caribæa 432.
Machaonia cymosa 435.
Lindeniana 48.
Maclura tinctoria 358.
Macrophoma Surinamensis 285.
Macroscepis rotata 38.
Macrosiphonia Berlandieri 383.
Malachra alceæfolia 508.
capitata 378, 508.
palmata 307, 508.
triloba 307.
MALPIGHIACEÆ 24, 301, 369, 428, 497.
Malphighia coccigera 428.
glabra 498.
glabra acuminata 400.
setosa 428.
undulata 24.
urens 428, 498.
urens lanceolata 498.
MALVACEÆ 29, 307, 377, 430, 507.
Malva spicata 29.

Malvastrum Coromandelianum 508.
Coromandelianum 378.
spicatum 508.
spicatum 29.
tricuspidatum 378.
Malvaviscus arboreus 30, 307, 378.
concinnus 30.
Sepium 30.
Malveopsis spicata 29.
Mammea Americana 31, 511.
Mangifera Indica 32, 375, 504.
Manihot Carthaginensis 374.
Manihot 375, 502.
rhomboidea 375.
MARANTACEÆ 481.
Maranta Indica 481.
Mariscus capitlaris 476.
flavus 476.
rufus 476.
Marsdenia floribunda 537.
maculata 39, 314.
Mexicana 314.
umbellata 433.
MARTYNIACEÆ 46, 319, 391, 529.
Martynia diandra 46, 319, 391, 529.
Mastigolejunea auriculata 7.
Mathiola scabra 48.
Maximowiczia tripartita 322.
Maximiliana hibiscoides 31, 311, 380.
Mayepea Caribæa 518.
Maytenus elæodendroides 505.
lævigatus 505.
phyllanthoides 375.
Mogiphanes straminea 16.
Meibomia adscendens 22.
affinis 22.
albiflora 398.
axillaris 495.
incana 495.
mollis 495.
Neo-Mexicana 399.
scorpiurus 299, 368, 428, 495.
spiralis 495.
tortuosa 299, 495.
triflora pilosa 495.
Melampodium divaricatum 53, 324, 395.
hispidum 324.
paludosum 395, 535.
MELAMSPORACEÆ 465.
Melanthera deltoidea 52, 323.
hastata 395.
hastata 52, 323.
MELASTOMACEÆ 36, 431, 516.
Melastoma Xalapensis 36.
MELIACEÆ 26, 302, 370, 500.
Melia Azedarach 26, 302, 500.
Melicocca bijuga 505.
oliviformis 403.
Melocactus communis 513.
Melochia hirsuta 430.
nodiflora 510.
pyramidata 31, 510.
tomentosa 31, 310, 379, 510.
Melothria fluminensis 393.
pendula 50, 322, 393.
pervaga 532.
MENISPERMACEÆ 297, 362, 427, 488.
Mentha aquatica 525.
Mertensia lævigata 399.
Mesosphærum actinocephalum 434.
capitatum 525.
pectinatum 525.
suaveolens 525.
Metastelma hamatum 433.
parviflorum 38.
Schlechtendalii 314, 384, 519.
Miconia androsæmifolia 432.
angustifolia 516.
argyrophylla 516.
impetiolaris 516.
lævigata 516.
præcox 431.
prasina 431.
Wrightii 431.
Micromeria Brownei 43, 317.
Microspora amœna 286.
Microtea debilis 486.
Mikania cordifolia 395.
gonoclada 536.
Houstonis 51.
scandens 324.
Milium compressum 471.
punctatum 471.
Milleria quinquefolia 395.
Mimosa asperata 368.
Ceratonia 492.
cornigera 19.
Farnesiana 19.
glauca 19.
pudica 492.
somnians 368.
trijuga 20.
Mirabilis Jalapa 296, 361, 486.
longiflora 16.
Mollugo nudicaulis 487.
Momordica Charantia 50, 394, 533.
cylindrica 50.
Monanthochlöe littoralis 288.
Monniera Dianthera 527.
Monniera 44, 390, 528.
procumbens 44, 319, 390.
Montanoa grandiflora 395.
hibiscifolia 324.
MORACEÆ 14, 293, 358, 482.
Morinda citrifolia 531.
Royoc 49, 321, 392, 435.
MORINGACEÆ 298, 362, 490.
Moringa aptera 298, 362.
Moringa 490.
Morrisonia Americana 490.
Morus celtidifolia 293, 359.
tinctoria 358.
Mouriria Domingensis 516.
Mucuna pruriens 368, 496.
Muntingia Calabura 377.
Murraya exotica 499.
exotica 25.
MUSACEÆ 13, 358, 480.
Musa paradisiaca 13, 480.
sapientum 13, 358, 480.
MUSCI 7, 286, 347.
MYCOSPHÆRELLACEÆ 465.
Myginda Gaumeri 401.
latifolia 505.
pallens 32, 505.
pallens 32.
Myrica coriacea 515.
coriacea Imrayana 515.
Myrodia turbinata 510.
Myrospermum frutescens 493.
Myroxylon nitidum 32.
Peruiferum 368.
MYRSINACEÆ 36, 312, 382, 517.
MYRTACEÆ 36, 312, 381, 431, 514.
Myrtus Pimenta 36, 381.

Nama Jamaicensis 40, 315, 521.
Nasturtium officinale 489.
Naudinia argyrophylla 431.
Nazia aliena 471.
Nectandra Antillana 489.
exaltata 427.
membranacea 489.
patens 427.
Willdenoviana 18.
Nephrolepis acuta 469.
exaltata 468.
Nerium Oleander 383, 518.
Neurolæna lobata 54.
Nicotiana glutinosa 388.
pilosa 388.
Tabacum 527.
NIDULARIACEÆ 467.
Nostoc verrucosum 347.
NYCTAGINACEÆ 17, 296, 361, 427, 486.
NYMPHÆACEÆ 17, 296, 361, 488.
Nymphæa ampla 17.

Ocimum Basilicum 318, 387, 525.
canum 318.
micranthum 318, 387, 525.
Selloi 318.
Octoblepharum albidum 286.
Odontonema nitidum 530.
Œnothera octovalvis 516.
OLACACEÆ 294, 484.
Oldenlandia callitrichoides 531.

Oldenlands corymbosa 434, 531.
OLEACEÆ 518.
Oliganthes oxylepis 51.
ONAGRACEÆ 312, 516.
Oplismenus Burmannii 352.
hirtellus 473.
setarius 473.
Opuntia coccinellifera 513.
Curassavica 513.
horrida 513.
spinosissima 513.
tuberculata 513.
Tuna 35, 513.
ORCHIDACEÆ 13, 292, 426, 481.
Orthopogon Burmannii 352.
hirtellus 473.
Ossæa macrandra 432.
OXALIDACEÆ 300, 369, 428, 497.
Oxalis Berlandieri 300, 369.
corniculata 497.
corniculata microphylla 497.
frutescens 428.
latifolia 301, 369, 497.
Martiana 497.
Oxandra laurifolia 488.
Oxybaphus violacea 296.

Pachira alba 309.
fastuosa 379.
fastuosa 309.
macrocarpa 30.
Pachyanthus Cubensis 432.
Pachyrhizus angulatus 300.
Pachystachys coccinea 537.
Padina pavonia 468.
Pæpalanthus androsaceus 426.
Palicourea Pavetta rosea 531.
PALMACEÆ 11, 290, 354, 426.
PANDANACEÆ 9.
Pandanus candelabrum 9.
utilis 9.
Panicum bambusoides 472.
barbinode 472.
brevifolium 288.
cæspitosum 473.
Carthaginense 353.
caudatum 473.
colonum 288, 353, 472.
crus-galli 353.
Dactylon 474.
diffusum 472.
divaricatum 288, 353, 472.
divaricatum latifolium 353.
divaricatum pubescens 472.
fuscum 288, 353, 472.
fuscum fasciculatum 472.
glaucum 473.
glutinosum 472.
glutinosum Lam. 472.
grossarium 472.
Panicum insulare 288.
insulare 353.
insularis 473.
jumetorum 353, 472.
lanatum 353.
lanatum 353.
latifolium 472.
leucophæum 353.
Lindeni 9.
Linkianum 353.
marginatum 353.
maximum 353, 472.
molle 472.
paspaloides 473.
plantagineum 473.
procumbens 473.
prostratum 473.
prostratum pilosa 473.
sanguinale 9, 288, 353.
sanguinale 473.
setosum 473.
PAPAVERACEÆ 18, 297, 362, 489.
Parathesis crenulata 37.
Parietaria Pennsylvanica 293.
Paritium tiliaceum 508.
Parkinsonia aculeata 493.
Parmentiera cereifera 390.
edulis 45, 390.
Parosela Domingensis 21, 299, 368.
Parthenium Hysterophorus 52, 324, 535.
Paspalum cæspitosum 471.
ciliatum Lam. 471.
compressum 471.
conjugatum 471.
distichum 471.
elongatum 353.
fimbriatum 472.
glabrum 472.
lentiginosum 353.
paniculatum 288.
paspaloides 354.
platycaule 471.
plicatum 472.
vaginatum 354, 472.
virgatum 472.
PASSIFLORACEÆ 34, 311, 380, 512.
Passiflora Andersonii 34.
ciliata 311.
ciliata 380.
fœtida 34, 380, 512.
gossypiifolia 311.
hirsuta 512.
incarnata 512.
laurifolia 512.
pallida 512.
pulchella 380.
quadrangularis 512.
rubra 512.
suberosa 380, 512.
Pauletia inermis 364.
Paullinia Curassavica 33.
Curassavica 402.
Cururu 402.
frutescens glabrescens 376, 402, 505.
Mexicana 33.
Pavonia racemosa 430, 508.
spinifex 508.
Pectis elongata **Schottii** 396.
humifusa 535.
linifolia 396.
pectinata 396.
prostrata 324, 396.
punctata 325, 535.
Pedilanthus angustifolius 504.
Itzæus 305, 375.
nodiflorus 305.
padifolius 504.
tithymaloides 504.
Pellæa aspera 287.
Pencillium glaucum 286.
Pencillius capitatus 467.
Peperomia acuminata 481.
alata 538.
Cubana 481.
glabella 482.
glutinosa 293.
Guadeloupensis 538.
Langsdorffii 538.
myrtifolia 538.
obtusifolia 482.
pellucida 482.
polystachya 482.
Pera oppositifolia 429.
Perdicium radialis 54.
Pereskia aculeata 513.
Bleo 513.
PERISPORACEÆ 465.
PERONOSPORACEÆ 465.
Persea gratissima 17, 297, 362.
Persea 489.
Pestalozzia **Coccolobæ** 286.
Petitia Domingensis 524.
Pœppigii 434.
Petiveria alliacea 16, 295, 360, 486.
Petræa arborea 317, 387.
volubilis 42, 524.
Peucedanum graveolens 517.
Pharbitis acuminata 521.
cathartica 521.
Nil 521.
purpurea 521.
Pharus glaber 474.
lancifolius 474.
Phaseolus alatus 497.
dysophyllus 368.
lunatus 23, 300, 368, 497.
semierectus 497.
Truxillensis 23.
vulgaris 23, 497.
Phegopteris rudis 349.
tetragona 349.

Philibertia Evendbergii 314.
Lindeniana 38.
viminalis 433.
Philodendron lacerum 11.
trifoliatum 355.
Philoxerus vermicularis 295, 360.
Phœbe Antillana 489.
Phœnix dactylifera 355.
Phoradendron flavens 484.
flavescens 294.
Martinicense 484.
Phthirusa Caribæa 483.
Phyllachora graminis 465.
Phyllanthus acuminatus 26, 306.
Carolinensis 306.
cyclanthera 429.
diffusus 429.
distichus 500.
Niruri 26, 501.
nobilis 501.
nobilis Antillana 501.
Physalis angulata 388, 434, 527.
Barbadensis obscura 318.
fœtens 388.
mollis cinerascens 318.
pubescens 388, 527.
Physalospora immorata 465.
Sp. 465.
PHYTOLACCACEÆ 16, 295, 360, 486.
Phytolacca Mexicana 295.
octandra 295.
Picræna excelsa 500.
Picramnia antidesma 25.
Pictetia aculeata 494.
Pilea muscosa 294.
succulenta 483.
Pimenta acris 515.
officinalis 381.
vulgaris 36, 515.
vulgaris 381.
Pimpinella Anisum 517.
PINACEÆ 425.
Pinus Cubensis 425.
PIPERACEÆ 13, 293, 358, 427, 481, 537.
Piper Amalago 537.
auritium 358.
Bredemeyeri 481.
incurvum? 537.
medium 291, 481.
peltatum 481.
tuberculatum 427.
Yucatanense 13.
Piscidia Erythrina 368, 428.
Erythrina 22, 300.
Pisonia aculeata 296, 361, 487.
inermis 487.
rotundata 427.
subcordata 487.
Pitcairnia angustifolia 478.
Pithecoctenium hexagonum 391.
Pithecolobium brevifolium 300.
ligustrinum 19.
oblongum 19.
Saman 491.
sericifolium 19.
unguis-Cati 19, 300, 491.
unguis-Cati ferfex 491.
Pittiera longipedunculata 393.
PLANTAGINACEÆ 47, 321, 392, 530.
Plantago major 47, 321, 392.
major tropica 530.
PLEOSPORACEÆ 465.
Pleroma longifolia 36.
Pluchea camphorata 396, 535.
odorata 52, 325, 396, 535.
purpurascens 52, 396, 535.
PLUMBAGINACEÆ 37, 313, 382, 517.
Plumbago Capensis 382.
scandens 37, 313, 382, 517.
Plumeria alba 38, 518.
obtusifolia 518.
pudica 383.
rubra 313, 383, 519.
Poa ciliaris 475.
Podopterus Mexicanus 294, 359.
Poinciana pulcherrima 21.
regia 300, 493.
Polanisia icosandra 490.
POLYGALACEÆ 26, 428, 500.
Polygala paniculata 26.
uncinata 429.
POLYGONACEÆ 15, 294, 359, 427, 484, 537.
Polygonum glabrum 427.
Polyodon distichum 350.
POLYPODIACEÆ 468.
Polypodium aureum 469.
crenatum 469.
incanum 349, 470.
incisum 470.
lycopodiodes . 8, 287, 349.
molle 468.
Phyllitidis 8, 349, 470.
piloselloides 470.
reptans 349.
serpens 470.
Swartzii 470.
tetragonum 470.
Polystictus sanguineus 467.
PONTEDERIACEÆ 292, 479.
Porana paniculata 521.
Porophyllum Ervendbergii 396.
nummularium 325.
Portlandia grandiflora 531.
PORTULACACEÆ 296, 361, 487.
Portulaca halimoides 296, 488.
oleracea 296, 361, 487.
oleracea micrantha 487.
pilosa 296, 361, 487.
Portulaca quadrifida 487.
POTAMOGETONACEÆ 470.
Priva echinata 41, 317, 387.
lappulacea 41, 317, 387, 524.
Prosopis juliflora 368.
Protium heptaphyllum 370, 428.
Psidium Guajava 36, 312, 381, 515.
Psychotria Brownei 531.
glabrata 531.
horizontalis 531.
tenuifolia 531.
tubulosa 435.
undata 392.
Ptelea pentandra 25.
Pteris aquilina caudata 8.
caudata 8.
longifolia 469.
Pterocarpus Draco 368.
Pterocaulon virgatum 535.
PUCCINIACEÆ 466.
Puccinia convolvuli 466.
heterospora 466.
menthæ 466.
opulenta 466.
spermacosis 466.
vernoniæ 466.
PUNICACEÆ 312, 381, 514.
Punica Granatum 312, 381, 514.

Quamoclit coccinea 521.
Quamoclit 521.
Quararibea **Fieldii** 309, 379.
Quassia amara 499.
Quisqualis Indica 516.

Rachicallis Americana 47.
rupestris 47.
Ramalina calicaris farinacea 347.
calicaris fraxinea 347.
rigida 347.
Randia aculeata 47, 392, 531.
aculeata mitis 532.
latifolia 47.
longiloba 47.
Mussænda 532.
Xalapensis 321.
RANUNCULACEÆ 17, 297, 362.
Raphanus Raphanistrum 362.
Rauwolfiia heterophylla 37, 313, 383.
Lamarckii 519.
nitida 519.
Renealmia sylvestris 480.
RESTIONACEÆ 426.
Reynosia mucronata 506.
RHAMNACEÆ 33, 376, 506.
Rhexia longifolia 36.
RHIZOPHORACEÆ 35, 312, 381, 514.
Rhizophora Mangle 35, 312, 381, 514.

Rhodomelaceæ 468.
Rhodophyllidaceæ 468.
Rhœo discolor 292, 356, 479.
Rhus Antillana 504.
Metopium 32.
Rhynchosia minima 23, 368.
minima lutea 496.
Rhynchospora micrantha 354.
scutellata 425.
Ricinus communis 27, 306, 375, 502.
Rivina humilis 16, 486.
humilis glabra 16, 295.
humilis scandens 16.
lævis 16.
scandens 16.
Rochefortia acanthophora 523.
Rondeletia correifolia 434.
pilosa 532.
stenosiphon 47.
Rosaceæ 19, 427, 490.
Rosmarinus officinalis 387.
Roulinia Jacquinii 38.
Rubiaceæ 47, 321, 392, 434, 530.
Ruellia albicaulis 320, 391.
coccinea 529.
geminiflora 391.
paniculata 46, 320.
strepens 529.
tuberosa 320, 391, 529.
Tweedyana 320.
Ruppia rostellata 470.
Russelia juncea 319, 528.
sarmentosa 44, 319.
Rutaceæ 25, 301, 370, 401, 499.
Ruta graveolens 301.

Sabal Blackburnianum 426.
Mexicanum 11, 355.
Sabinea florida 494.
Saccharum officinarum 471.
Sachsia polycephala 435.
Sagræa macrandra 432.
Salacia **Blainii** 430.
Salicaceæ 482.
Salix Humboldtiana 482.
Salicornia ambigua 485.
Bigelovii 360.
mucronata 360.
Salvia coccinea 43, 318, 387, 525.
coccinea ciliata 525.
occidentalis 43, 318, 526.
serotina 43, 526.
Salviniaceæ 349.
Salvinia auriculata 349.
Sambucus Canadensis 321, 393.
Mexicana 49.
Samyda grandiflora 431.
serrulata 389, 512.
Sansevieria Guineensis 479.
Sanvitalia procumbens 325, 396.
Sapindaceæ 33, 306, 376, 402, 505.
Sapindus inæqualis 506.
saponaria inæqualis 402.
Sapotaceæ 37, 313, 382, 517.
Sarcostemma Lindenianum 38.
Sargassum ludigerum 467.
Savia sessiliflora 501.
Scævola Plumieri 50, 394, 533.
Schæfferia frutescens 505.
Schmidelia comina 33.
Cominia 402.
occidentalis 506.
Schœnus capillaris 476.
Schœpfia arborescens 484.
Schwenkia Americana 319, 389.
Scirpus articulatus 537.
capitatus 477.
interstinctus 477.
lacustris 290.
Scleria lithosperma 477.
pterota 477.
Sclerocarpus divaricatus 396.
Scleropus amarantoides 486.
Scolosanthus versicolor 532.
Scoparia dulcis 528.
Scrophulariaceæ 44, 319, 389, 434, 527.
Sebastiania corniculata 429.
Securidaca Brownii 500.
erecta 500.
Securinega Acidothamnus 501.
Selaginella **longispicata** 287, 349.
Senecio Berlandieri 396.
salignis 325.
Seriana plicata 33.
Serjania **adiantoides** 403.
atrolineata 403.
goniocarpa 403.
lucida 505.
Mexicana 33.
plicata 33, 403.
polyphylla 376.
scatans 33.
scatans 403.
Sesbania grandiflora 368.
macrocarpa 369.
Sesuvium portulacastrum 296, 361, 487.
Setaria auriculata 289.
flava 288.
glauca 473.
latiglumis 289.
pauciseta 289.
setosa 473.
setosa caudata 473.
Setariopsis auriculata 289, 354.
latiglumis 289.
Sicydium tamnifolium 50.
tamnifolium 394.
Sida acerifolia 29.
acuta carpinifolia 29, 307, 378.
Sida angustifolium 508.
anomala 378.
arguta 508.
carpinifolia acuta 508.
carpinifolia Antillana 508.
carpinifolia brevicuspidata 509.
ciliaris 307, 509.
cordifolia 430, 509.
cordifolia althæfolia 509.
diffusa 308.
glutinosa 509.
hastata 29.
hederæfolia 430.
Indica 29.
nervosa viscosa 509.
pilosa 378.
retusa 509.
rhombifolia 308, 509.
rhombifolia retusa 509.
spinosa 509.
spinosa polycarpa 509.
supina glabra 509.
supina pilosa 308, 378, 509.
Sideroxylon Mastichodendron 313, 518.
Silene Cucubalus 296.
Silphium trilobatum 53.
Simarubaceæ 25, 301, 370, 499.
Simaruba glauca 370.
Sinapis arvensis 489.
brassicata 489.
Sinebra repens 350.
Siphonoglossa sessilis 320, 392, 529.
Sisyrinchium sp. 426.
Smilaceæ 12.
Smilax **Gaumeri** 357.
Havanensis 426, 479.
Mexicana var. 12.
Solanaceæ 43, 318, 388, 434, 526.
Solanum aculeatissimum 527.
Amazonicum 44, 388.
callicarpæfolium 388.
fuscatum 388.
Havanense 44.
Humboldtii 44.
igneum 527.
Jamaicense 389.
Juripeba 389.
mammosum 389, 527.
Melongena 389, 527.
nigrum 44.
racemosum 527.
rostratum 389.
Seaforthianum 318, 527.
torvum 44, 318, 389, 527.
verbascifolium 44, 318, 389, 527.
Sonchus oleraceus 325, 535.
Sophora tomentosa 21, 493.
Sorghum vulgare 471.

Sparganophorus Vaillantii 535.
Spermacoce Hænkiana 49.
Hænkiana 321.
lævis 49.
podocephala 435.
tenuior 49, 321, 393, 532.
tenuior angustifolia 532.
verticillata 49, 321, 393.
Sphærococcaceæ 468.
Spigelia **Blainii** 432.
Spilanthes Beccabunga 53.
repens 53.
Spiranthes orchioides 292.
polyantha 293.
Spironema fragrans 356.
Spondias dulcis 504.
lutea 32, 375, 504.
purpurea 375, 504.
Sporobolus Domingensis 474.
Indicus 474.
littoralis 474.
minutiflorus 290, 474.
Virginicus 474.
Stachytarpheta Jamaicense 41, 317, 525.
Staphyleaceæ 430.
Stemmadenia insignis 383.
Stenandrium droseroides 434.
Stenostomum lucidum 530.
Stenotaphrum Americanum 10.
Americanum 474.
secundum 474.
Stephanomeria runcinata 325.
Sterculiaceæ 31, 379, 430, 510.
Sterculia Carthaginensis 31, 379.
tereophyllum leucostegum 7.
perpusillum 348.
Stigmatophyllon lupulus 369.
periplocæfolium 498.
Streptachne tenuis 354.
Strumpfia maritima 48.
Stylosanthes biflora 369.
hamata 495.
viscosa? 495.
Suæda fruticosa 360.
Suriana maritima 25, 301, 499.
Swietenia Mahogani 500.
Symplocaceæ 432.
Symplocos Martinicense 432.
salicifolia 432.
Synedrella nodiflora 54, 397.
nodiflora 535.
Syntherisma insularis 473.
sanguinalis 473.
setigera 473.

Tabernæmontana Acapulcensis 38, 383.
amygdalæfolia 38, 313.
Tænitis lanceolata 469.
Tagetes patula 325, 397.
Tagetes tenuifolia 54.
Talinum patens 487.
triangulare 296, 487.
Talisia olivæformis 403.
Tamarindus Indica 20, 300, 369, 492.
Tamaricaceæ 511.
Tamarix Indica 511.
Tamonea scabra 317.
Tecoma Capensis 528.
lepidophylla 434.
leucoxylon 528.
Stans 45, 319, 391, 529.
Tephrosia cinerea 22, 299.
Teramnus labialis 496.
uncinatus 496.
uncinatus albiflorus 496.
Terentepohlia rigidula 467.
Terminalia Catappa 36, 312, 381.
Ternstrœmia elliptica 511.
Tetramerium costatum 47, 320.
costatum 391.
hispidum 47, 321.
Tetrapteris inæqualis 370.
Tetrazygia argyrophylla 431.
elæagnoides 516.
Teucrium inflatum 43, 318.
Thalassia testudinum 470.
Theaceæ 430, 511.
Theobroma Cacao 31, 510.
Thespesia populnea 509.
Thevetia Gaumeri 38.
nereifolia 38, 313, 383, 519.
nitida 38.
spathulata 383.
Thouinia **paucidentata** 403.
Thrinax argentea 355.
parviflora 11.
Thuidium involvens 286.
Thunbergia alata 537.
fragrans 530.
volubilis 530.
Thymelæaceæ 431.
Thymus Brownei 43.
Tibouchina longifolia 36.
Tigridia violacea 13.
Tiliaceæ 34, 307, 376, 430, 506.
Tillandsia Balbisiana 291.
brachycaulos 356.
circinnata 291.
dasyliriifolia 291.
fasciculata 291.
fasciculata latispica 291, 356.
filifolia 12, 291.
polystachya 12, 291.
recurvata 291, 356, 478.
setacea 356.
streptophylla 291, 356.
usneoides 292, 478.
utriculata 292, 356, 478.
Tillandsia vestita 12, 291, 356.
Tithonia diversifolia 325, 397.
tagetiflora 397.
tubæformis 325.
Tobinia punctata 499.
Torulinium Eggersi 476.
Michauxianum 476.
Tournefortia fœtidissima 523.
gnaphalodes 18, 41, 316, 386, 523.
hirsutissima 523.
volubilis 41, 315, 433, 523.
volubilis microcarpa 523.
Tragia nepetæfolia 306, 375.
volubilis 502.
Tragus occidentalis 471.
Trema micrantha 482.
Trianosperma graciliflora 533.
Trianthema monogyna 487.
Tribulus cistoides 498.
cistoides 24.
maximus 24, 301, 370.
terrestris cistoides, 24, 301, 370.
Triceratia bryonoides 394.
Trichilia hirta 500.
spondioides 370.
Terminalia 26, 302.
Tricholæna insularis 473.
Tri-uspis latifolium 352.
Tridax procumbens 54, 325.
procumbens pubescens 325, 397.
Trilix crucis 511.
Triodon angulatum 49.
Triphasia trifoliata 499.
Triumfetta althæoides 34, 506.
Lappula 506.
rhomboidea 507.
semitriloba 380, 506.
semitriloba Havanensis 507.
Trixis frutescens 397.
frutrescens 54.
Inula 54.
radialis 54, 326.
radialis 397.
Tubiflora squamosa 46, 320, 392.
Turneraceæ 311, 380, 512.
Turnera diffusa aphrodisiaca 311, 380.
ulmifolia 311, 512.
ulmifolia acuta 512.
Typhaceæ 470.
Typha angustifolia Domingensis 470.
Domingensis 470.

Ucacoua nodiflora 535.
Udotea flabellata 467.
Ulmaceæ 399, 482.
Umbelliferæ 312, 381, 517.

Uniola spicata 351.
Uredo commelinacea 466.
gouanæ 466.
Urena Americana 509.
Urera baccifera 359.
elata 483.
microcarpa 14, 294, 359.
Uromyces Howei 466.
URTICACEÆ 14, 293, 359, 483.
Urvillea ulmacea 307.
ulmacea genuina 403.
USTILAGINACEÆ 466.

Valerianodes Jamaicensis 41, 317, 392, 525.
Vandellia diffusa 528.
Vanilla anaromatica 426.
Varronia globosa 40.
VERBENACEÆ 41, 316, 386, 434, 523.
Verbena Jamaicensis 41.
lappulacea 41.
nodiflora 42.
prismatica 41.
squamosa 46, 320.
stœchadifolia 42.
Verbesina alata 535.
dichotoma 52.
Vernonia arborescens 535.
arborescens divaricata 535.
arborescens Lessingiana 536.
arborescens Swartziana 535.
lanceolaris 50.
Schiedeana 50.
Vigna Catjang 497.
repens 497.
Virguiera helianthoides 53, 326, 397.
Vilfa Domingensis 474.
Villamilla octandra 486.
octandra 16.
Vinca glabra 37.
rosea 313, 384, 519.
VIOLACEÆ 32, 311, 404, 511.
Viola hybanthus 32.
VITACEÆ 33, 307, 506.
Vitis arborea 307.
bifinnata 307.
Caribæa 506.
rhombifolia 307.
sicyoides monstrosa 33.
tiliæfolia 33.
Vitex divaricatus 525.
pyramidata 317.
Volkameria ligustrina 42.
Vouacapoua Americana 495.
Vriesia psittacina decolor 292.

Waltheria Americana 31, 380.
Indica 310, 511.
Wedelia buphthalmoides 536.
buphthalmoides Antiguensis 536.
buphthalmoides Dominicensis 536.
carnosa 53.
cruciana 536.
hispida 326.
trilobata 53, 536.
Willughbæa cordifolia 536.
Willughbæa gonoclada 536.
Houstonis 51.
Wissadula mucronulata 378.
periplocæfolium 510.
tricarpellata 378.
Wolffia Braziliensis 290, 355.

Xanthium Canadense 397.
macrocarpum 536.
Strumarium 536.
Xanthosoma atrovirens 478.
hastatum 478.
sagittæfolium 478.
Xanthoxylum Clava-Herculis 499.
flavum 499.
Xiphidium floribundum 426.
Xylosma nitidum 32.

Yucca gloriosa 479.
Yucatana 12, 357.

Zanthoxylum pterota 24, 370.
Zebrina pendula 356, 479.
Zephyranthes Lindleyana 357.
rosea 479.
Zexmenia Costaricensis 397.
hispida 326.
trachylepis 53.
ZINGIBERACEÆ 426, 480.
Zingiber officinale 426, 480.
Zinnia elegans 536.
Zizyphus reticulata 506.
Zornia diphylla 428.
ZYGOPHYLLACEÆ 24, 301, 370, 428, 498.

LIBRARY
OF THE
UNIVERSITY OF ILLINOIS

FICUS RUBIGINOSA?

FIELD COLUMBIAN MUSEUM

PUBLICATION 4.

Botanical Series. Vol. I, No. I.

CONTRIBUTION

TO THE

FLORA OF YUCATAN

BY

CHARLES FREDERICK MILLSPAUGH,

Curator, Department of Botany.

CHICAGO, U. S. A.

August, 1895.

Contribution to the Flora of Yucatan.

The peninsula of Yucatan forms the southeastern extremity of Mexico, and constitutes its two Confederate States of Yucatan in the east and Campeche in the west. It projects as a compact rectangular mass between the waters of the Gulf of Mexico on the north and those of the Caribbean Sea on the south, and is separated from the Island of Cuba on the east by the narrow channel of Yucatan, through which pours the current of the Gulf Stream, after being heated in the cauldron of the southern portion of the Gulf. The coast line of the peninsula, about 700 miles in extent, presents a uniform, monotonous aspect of an almost continuous line of low sand dunes, which enclose at nearly all points broad lagoons, ranging in depth of water according to the season. The peninsula has a mean breadth of about 200 miles, and a total area of 55,400 square miles, being about the size of the State of Illinois,

The entire eastern portion of the country is, geologically speaking, of recent formation, having been built up in shallow water by polypi, in the form of a low level plateau of porous limestone, having only sufficient soil caught in its uneven surface to support a dense but dwarfish vegetation.

From Cape Catoche, its most northeasterly projection, southward along the coast, extend the great Blanquilla reefs, among which are scattered many small islands of coralline rock and clear coral sand. The principal of these are Cankun, Holbox, Cantoy and Mugeres; all lie near the coast, and their formation is apparently too recent to create much soil except the sand of the disintegrated reef rock. About half way down the coast, and ten miles from it in the channel, there is, however, a larger and older island, Cozumel, which is doubtless a special formation, being considerably higher and older than any of the others. This island has a soil similar to, if not the same as, that of the mainland, and its vegetation is less of the sand-littoral than that of the others. Cozumel is about twenty-five miles long, its highest point being nearly one hundred feet above the sea. In speaking of the flora, Professor Hemsley says in the supplement to

Vol. IV, of Godman and Salvin's *Biologia Centrali-Americana*: "Taking the present data, the flora of Cozumel is probably more West Indian than continental American in character; for, although the number of species, excluding doubtful ones common to Cozumel and the Continent, but not found in the West Indies, is somewhat larger than the number of West Indian species found in Cozumel, but not on the mainland, those of the latter category are of a more distinct type than those of the former." This may prove true, yet an element enters into the collection upon which Professor Hemsley based his observation of which he had no knowledge; the collector, Dr. Gaumer, a resident of the mainland, collected on Cozumel only such species as he had not observed on the Peninsula.

This section of Mexico is, in the main, excessively dry, there being no elevations sufficiently high to intercept the moisture-bearing clouds from the Atlantic, while those from the Pacific are shut out by the Sierra Madre range; rainfall is deficient, especially in the northern and southern districts. The heat is also great here, and almost continuous through the year. A good idea of the climate is given in the following letter recently received from Dr. Gaumer: "March is one of our dryest months, quite hot but very healthy; many of the forest trees are then in bloom, though nearly leafless; for hunting wild animals and birds it is, however, the best month in the year. January and February are not so hot and have more flowers, while April and May are much hotter and the flowers are fewer still. In October, November and December, flowers are very abundant and the weather cool. June is by far the most delightful month of all the year; Nature, clad in the splendor of her fresh green attire, is everywhere bedecked with beautiful flowers, among which flit myriads of bright-hued butterflies and brilliant plumaged birds; rains are frequent, however, and form the only drawback to collecting. July, August and September are unsafe months for foreigners to visit this region, although there is no danger if they come in January and remain through this period."

Although the upper and main portion of Mexico has been for years a favorite collecting ground for botanists, Yucatan has, strangely enough, received but little attention, and so little is known of its flora that Drude in his *Handbuch der Pflanzengeographie*, says nothing whatever of the region, though the work was published in 1890. Grisebach, who is best equipped concerning the flora of the neighboring Gulf islands, simply states in his *Vegetation der Erde* (1884) that the conditions of the region prevent luxuriance of vegetable growth, and Hemsley in his report on the botany of Mexico and Central America prepared for the *Biologia* quoted above, indicates the paucity of

knowledge concerning the plant life of the peninsula in the following words: "We may add that little is known of the details of the botany of Yucatan, except that it is very poor and scanty, and largely composed of plants that still bear long droughts without injury. The poverty of the flora is ascribed to the fact that the copious rains rapidly filter away through the porous limestone substratum." All this renders the botany of the country of still greater interest, and its development of the greatest. It is, therefore, for this reason that the Museum has decided to put a collector at work in the field and accumulate full knowledge of its flora. In order, therefore, that our work may be as complete as possible, we have carefully compiled in this first contribution all that we could find published concerning previous collections.

The history of previous work toward a knowledge of this region is that of fragmentary collections, no collector having so far done any exhaustive collecting in the field.

In 1835, Jean Jules Linden, a Belgian horticulturist, gathered about twenty-five species of plants in Yucatan while on his way to Vera Cruz. This constitutes the first known collection from this part of Mexico.

In 1848, Hon. E. P. Johnson collected about seventy-five species, which he sent to Dr. Torrey with the label "Yucatan and Tabasco." Some doubt is therefore felt as to the purely Yucatan specimens in the gathering.

In 1885 and 1886, Dr. George F. Gaumer, while collecting in ornithology for the British Museum, gathered on the islands of the east coast, the inflorescence and a few leaves of by far the largest number of plants that have come out of the country, 224 species. These, together with the two collections mentioned above, were worked over by Prof. Hemsley for the *Biologia*, and comprise all that is known of the insular flora of Yucatan.

The collection which forms the basis of this contribution was made last January, during an expedition to the ruined city of Chichen Itza, and the Islands of Mugeres and Cozumel, and is again incomplete and fragmentary, on account of the excessive dryness of the season, and the rapidity of movement of the party. The expedition was generously planned and carried forward by Mr. Allison V. Armour of Chicago, and as it was conducted in his steam yacht it gave an opportunity of visiting the islands, that would otherwise have been unattainable.

In the following list, the species in black face type are those collected or observed by myself, the balance, in the body type, have been carefully compiled from the pages of the *Biologia Centrali-*

Americana, as previously noted. The arrangement of the natural orders is that of Engler and Prantl in *Die Naturlichen Pflanzenfamilien*, as far as published, otherwise that of Bentham and Hooker's *Genera Plantarum*. The nomenclature is in accordance with the "Rochester rules" as far as our reference facilities would permit.

The determination of the species was made by the author in the Herbarium of the Shaw Botanical Gardens at St. Louis, through the courtesy of the Director, Prof. Wm. Trelease, with the exception of a few which were kindly compared in the Gray Herbarium, Harvard, by Prof. B. L. Robinson, and the orders credited to specialists in the body of the list.

CHICAGO, August, 1895.

Plants of the Allison V. Armour Expedition.*

HEPATICAE.†

MASTIGOLEJEUNEA, Spruce.

M. auriculata, Wils.
On trunks of smooth trees. Chichen Itza, Jan. 17, 1895 (**106**).

CHEILOLEJEUNEA, Spruce.

C. ——— (Juvenile).
On bole of living tree. Mascab Pixoy, Jan. 22, 1895 (**151**).

C. ——— (Juvenile).
On trunk living Ceibo. Island of Cozumel, Jan. 5, 1895 (**37**).

EULEJEUNEA, Spruce.

E. lepida, L. & G.
On bark of living Ceibo. Island of Cozumel, Jan. 5, 1895 (**32**).

MUSCI.§

BARBULA, Hedw.

B. agraria, Hedw.
On dry limestone cobble, Cedral, Island of Cozumel, Jan. 5, 1895 (**48**). Associated with Didymodon æneus on dry limestone rock, Chal-tun-ha road, near Izamal, Jan. 1895 (**80** in part).

DIDYMODON, Hedw.

D. æneus, Sch.
Associated with Barbula agraria on dry limestone rock, Chal-tun-ha road, near Izamal, Jan. 14, 1895 (**80** in part) sterile.

STEREOPHYLLUM, Hedw.

S. leucostegum (Brid).
On limestone rock, Island of Cozumel, Jan. 5, 1895 (**33**). On dead and decaying limb, Lagoona Colombia, Island of Cozumel, Jan. 7, 1895 (**47**).

---

*In black face type

†Specimens kindly determined by Professor L. M. Underwood, who also referred the two Cheilolejeunea to Professor Stephani.

§Kindly determined by M. Professor Jules Cardot.

PTERIDOPHYTA.*

POLYDPODIUM, Linn.

P. Phyllitidis, Linn.

On fallen trunk of dead tree, Cedral, Island of Cozumel, Jan. 5, 1895 **(38)**.

Specimens very large and fine.

P. lycopodioides, Linn.

Climbing living trees in dry open woods, Chichen Itza, Jan. 18, 1895 **(134)**.

The same fine large form collected by Mr. Pringle in upper Mexico.

ADIANTUM, Linn.

A. tricholepis, Fee.

Mouth of small cave, north aspect of the Cerro Grande, Izamal, Jan. 13, 1895 **(54)**. Crevices of north aspect ruin of the Monjas, Chichen Itza, Jan. 17, 1895 **(112)**.

Both are excellent specimens of this once rare, beautiful fern.

PTERIS, Linn.

P. acquilina caudata (Linn.), Hook.

Pteris caudata, Linn. Collected by Dr. Gaumer on the Island of Cozumel, 1885.

HEMITELIA, R. Brown.

H. Hartii, Baker.

Collected by Dr. Gaumer on the Island of Cozumel, 1885.

CHEILANTHES, Swartz.

C. leucopoda, Link.

Collected by Dr. Gaumer on the Island of Cozumel, 1885.

ASPLENIUM, Linn.

A. pumilum, Swartz.

On sandy loam under limestone ledge, Cedral, Island of Cozumel, Jan. 5, 1895 **(35)**.

Fronds taller than is usual, the upper portion more elongated than in the type, and deeply pinnatifid to the acuminated apex, with the lobes more or less deeply cut, but not laciniated. Quite unlike in appearance the smaller normally deltoid forms with entire, or nearly entire upper third, but otherwise the characters are the same. The species, however, like all other Aspleneæ, is an extremely variable one, and the present form is no greatly marked exception.

Smaller plants with firmer texture collected at the same station on the under side of a small projecting limestone ledge **(36)**.

* Determinations and notes of my numbers kindly made by Mr. Geo. E. Davenport.

ASPIDIUM, Swartz.

A. trifoliatum, Swartz.
On the rocky walls of the Cenote de Cafetal, Chichen Itza, Jan. 18, 1895 (**132, 133**).
These two numbers embrace a series of forms from very small, young, and undeveloped plants upward, and prove especially interesting on this account.

ANEIMIA, Swartz.

A. adiantifolia, Swartz.
Mouth of small cave, north aspect of the Cerro Grande, Izamal, Jan. 13, 1895 (**58**).

CYCADACEAE.

DIOON, Lindl.

D. spinulosum, Dyer.
Progreso (C. *J. Hoege*).

PANDANACEAE.

PANDANUS, Linn. f.

P. utilis, L. f. **PAH.**
Margin of the Aguada of Xcholac, about two leagues south of Izamal.

P. Candelabrum, P. Beauv. **ƆIPIL.***
In the scrub between Progreso and the first railroad station south.

GRAMINEAE.

(*Maydeæ*).

COIX, Linn.

C. lachryma-Jobi, Linn. Job's Tears.
Collected in the Patio of the residence of Dr. Gaumer, Izamal, Jan. 13, 1895 (**73**).

(*Paniceæ*).

PANICUM, Linn.

P. sanguinale, Linn.
Collected by Dr. Gaumer on the Island of Cozumel, 1885.

P. Lindeni, Fourn.
Yucatan, type loc. ignot. (*Linden*).

CENCHRUS, Linn.

C. pallidus, Fourn.†
Crevices in rocks in the Patio of Dr. Gaumer, Izamal, Jan. 13, 1895 (**70**).
Note: Fourn. Mex. Pl. Gram. p. 50, = 465 Liebman from Santa Cruz.

* The inverted letter C is pronounced in the Maya *tz*.
† Kindly determined by Professor F. Lamson Scribner.

C. brevisetus, Fourn.*
Sands of the beach at Progreso, especially near the cattle pens where many cattle are herded on arrival from Tampico and other ports on the Gulf Coast of Mexico. Jan. 26, 1895 (**204**).
Note: Fourn. Mex. Pl. Enum. Gram. p. 50. = 3140 Bourgeau from Orizaba, Mexico.

STENOTAPHRUM, Trin.

S. Americanum, Schrank.*
Grassy edge of the sandy beach west shore, Island of Cozumel, at the portage to Lagoona Colombia, Jan. 7, 1895 (**50**).

(*Chlorideæ*).

CHLORIS, Sw.

C. Petræa, Linn.
Collected on the Island of Holbox by Dr. Gaumer, 1886.

BOUTELOUA, Lagasca.

B. Triæna (Spreng), Scribner.*
Top of pyramid of El Castillo, Chichen Itza, Jan. 19, 1895 (**135**).
Note: Scribner Proc. Acad. Nat. Sci. Phila., 1891, p. 307. *Antheropogon Triæna*, Spreng, Syst. 1, p. 293, ex. Kunth; *Bouteloua triathera*, Bentham (in part).

ELEUSINE, Gærtn.

E. Indica, Gærtn.
Collected on the Islands of Holbox and Cozumel by Dr. Gaumer in 1885-6.

E. Ægyptiaca, Pers.
Collected on the Island of Holbox by Dr. Gaumer, 1886.

Festuceæ.

ERAGROSTIS, Host.

E. ciliaris, Link.
Collected on the Islands of Cozumel, Mugeres and Holbox by Dr. Gaumer, 1885-6.

CYPERACEAE.

CYPERUS, Linn.

C. Brunneus, Sw.
Islands of Mugeres and Cozumel (*Gaumer*, 1886).

C. aristatus, Rottb.
Yucatan, loc. ignot. (*Schott*).

*Determined by Professor F. Lamson Scribner.

C. Ottonis, Boeckl.*
On the apex of the reef rocks, north end of the Island of Mugeres, Jan. 2, 1895 (**22**).

C. ligularis, Linn.
Island of Cozumel (*Gaumer*, 1885).

C. thrysiflorus, Jungh.
Island of Cozumel (*Gaumer*, 1885).

IRIA, Rich.

I, polymorpha (Boeckl.) O.K.
Fimbristylis polymorpha, Boeckl. Island of Holbox (*Gaumer*, 1886).

PALMAE.

THRINAX, Sw.

T. parviflora, Sw.
Throughout the Islands of Mugeres, Cankun and Cozumel. Island of Cozumel (*Gaumer*, 1885), "very abundant."

SABAL, Adans.

S. Mexicana, ? *Bayal.* **XAAN.**
The common palmetto of the Islands and the Main, growing often to a height of 80 feet. In woodlands everywhere.

S. ———— sp. (Hemsl.)
Island of Mugeres (*Gaumer*, 1886).

COCOS, Linn.

C. nucifera, Linn. *Coco.*
Islands of Contoy, Mugeres, Cancun and Cozumel, along the coast throughout, and as far inland as Izamal, though the tree does not thrive well at the latter place. The trees and fruits on Cozumel and Cancun are specially fine.

ARACEAE.

PHILODENDRON, Schott.

P. lacerum, (Jacq.) Schott?
Arum lacerum. Jacq. On the walls of the Cenote de Cafetal, Chichen Itza, in leaf only, Jan. 19, 1895.

BROMELIACEAE.

BROMELIA, Linn.

B. Pinguin, Linn. **CHOM**
Common along the bridle path from Mascab Pixoy to Chichen Itza, some specially fine specimens near Pisté.
Many specimens of Bromeliads and Orchids were seen on the trees of this region, but all were out of flower, and but few with old fruits, Jan. 17, 1895.

*Determined by Professor N. L. Britton.

ANANAS, Adans.

A. sativus, Mill. *Pina.* Pineapple.
The Pineapple is cultivated extensively on the Island of Cozumel, where the fruit is said to be the best produced in Mexico.

ÆCHMEA, R. & P.

A. laxiflora, Benth.
Yucatan, loc. ignot. (*Schott*).

HECHTIA, Kl.

H. Schottii, Baker.
Type locality, on the Cerro de Maxeana (*Schott*).

TILLANDSIA, Linn.

T. filifolia, Ch. & Schl.
Yucatan, loc. ignot. (*Johnson*).

T. polystachya, Linn.
Yucatan, loc. ignot. (*Johnson, Schott*).

T. vestita, Ch. & Schl.
Yucatan, loc. ignot (*Johnson*).

T. ——— sp. (Hemsl.)
Island of Holbox (*Gaumer*, 1886).

LILIACEAE.

YUCCA, Linn.

Y. Yucatana, Engelm.
Type locality, Ruins of Nohpat (*Schott*).

DASYLIRION, Zucc.

D. pliabile, Baker.
Yucatan, type loc. ignot. (*Schott*).

SMILACEAE.

SMILAX, Linn.

S. Mexicana, Kth. ? var., (Hemsl).
Island of Cozumel (*Gaumer*, 1885).

AMARYLLIDACEAE.

AMARYLLIS, Linn.

A. equestris (Herb.) Ait.
Hippeastrum equestre, Herb. Island of Cozumel (*Gaumer*, 1885).
Yucatan loc. ignot. (*Johnson*).

DIOSCOREACEA.

DIOSCOREA, Linn.

D. spiculiflora, Hemsl.
Yucatan type, loc. ignot. (*Linden*).

D. densiflora, Hemsl.
Island of Cozumel (*Gaumer*, 1886).

IRIDACEAE.

TIGRIDIA, Ker.

T. violacea, Schiede & Schl.
Yucatan, loc. ignot. (*Linden*).

MUSACEAE.

MUSA, Linn.

M. sapientum, Linn. Banana, **SAC HAAS.**
Cultivated on the Island and throughout the Main.

M. paradisiaca, Linn. Plantain, **BOX HAAS.**
Cultivated at San Miguel and Colombia, Island of Cozumel; at Merida, Izamal and Tuncas.

CANNACEAE.

CANNA, Linn.

C. Indica, Linn.
Island of Cozumel (*Gaumer*, 1886).

COMMELYNACEAE.

COMMELINA, Linn.

C. erecta, Linn.
Island of Cozumel (*Gaumer*, 1885).

C. Virginica, Linn.
Islands of Mugeres and Cozumel (*Gaumer*, 1886).

CALLISIA, Linn.

C. repens, Linn.
Yucatan, loc. ignot. (*Schott*).

ORCHIDACEAE.

CATTLEYA, Lindl.

C. ——— sp.
Abundant on various trees from Mascab Pixoy to Chichen Itza. In fruit only, Jan. 17, 1895.

PIPERACEAE.

PIPER, Linn.

P. Yucatanense, C. DC.
Yucatan, type loc. ignot. (*Linden*).

CASUARINACEAE.

CASUARINA, Linn.

C. Equisetæfolia, Linn.

Cultivated in the Campo Santo of Dolores, Island of Mugeres; Streets of San Miguel, Island of Cozumel; Island of Cankun; and in Merida, Izamal, and Progreso, Jan. 26, 1895 (**204**), height 30 to 60 feet.

MORACEAE.

FICUS, Linn.

F. laurifolia, Lam. *Laurel de la India.*

Cultivated extensively as an ornamental shade tree. Introduced from Cuba. Fine examples at Oncaan, Ouan, and Bocoba.

F. rubiginosa, ? (Plate I, Frontispiece). *Copo. Alamo.*

This form, to which we make doubtful reference in the absence of satisfactory flowers and fruits, is the most common species throughout the route we traveled on the Peninsula. It is a large tree 25 to 45 feet high, greatly spreading and very umbrageous. For some unaccountable reason it is called the *Alamo* (Poplar) by the Spaniards. Its juice contains sufficient caoutchouc to form, when mixed with that of the Sapote, an excellent bird lime.

F. grandifolia, ? **AKUM.**

A large tree 30 to 50 feet high. Fine specimens noted at Chichen Itza.

ALICASTRUM, P. Br.

A. Brownei, O.K. *Ramon.* **OX.**

Brosimum Alicastrum, Sw. Cultivated for fodder throughout the region embraced, also on the Islands of Mugeres and Cozumel. Fine specimens at Merida, Izamal, and Chichen Itza. Grows from 50 to 80 feet high. Yucatan, loc. ignot. (*Linden*).

The foliage "Ramon," serves as the principal green fodder for horses and mules from June to March. The wood is white and compact, and is used in carpentry; the fruit when boiled is eaten either alone or mixed with honey or corn meal, and the milky juice is used as an anti-spasmodic in the difficult breathing of asthma.

URTICACEAE.

URERA, Gaud.

U. microcarpa, Wedd.

Yucatan, loc. ignot. (*Johnson*).

ARISTOLOCHIACEAE.

ARISTOLOCHIA.

A. grandiflora, Sw.
Aristolochia foetens, Lindl. Cultivated at Izamal by Dr. Geo. F. Gaumer, who brought it from the Honduran Island of Ruatan in 1886. In flower, Izamal, Jan. 13, 1895 (**68**).

POLYGONACEAE.

ANTIGONON, Endl.

A. leptopus, Hook & Arn. ?
Island of Cozumel (*Gaumer,* 1885), "sepala exteriora basi subcordata," Hemsl.

COCCOLOBA, Linn.

C. Uvifera (Linn.) Jacq. *Uba.*
Polygonum Uvifera, Linn. On the inland edge of the sandy beach, Islands of Cozumel and Mugeres Jan. 1, 1895. Also collected here by Dr. Gaumer, 1885.

C. Cozumelensis, Hemsl.
Type locality, Island of Cozumel (*Gaumer,* 1885).

CHENOPODIACEAE.

ATRIPLEX, Linn.

A. cristata, H. B. K.
Island of Holbox (*Gaumer,* 1886).

LERCHEA, Hall.

L. ——— Sp. (Hemsl.)
L. Suaeda, Forsk. Island of Holbox (*Gaumer,* 1886).

AMARANTHACEAE.

CELOSIA, Linn.

C. paniculatata, Linn.
C. nitida, Vahl. Island of Mugeres (*Gaumer,* 1886).

AMARANTHUS, Linn.

A. hybridus, Linn. **XTEZ.**
A. chlorostachys, Willd. About the ruins of the old church of the Hacienda, Chichen Itza, Jan. 1895 (**129**).
The Maya name Xtez is applied to all similar Amaranths.

ACHYRANTHES, Linn.

A. aspera, Linn. *Sorillo blanco.* **SAC-PICHÉ.**
Waste grounds around the old church of the Hacienda, Chichen Itza, Jan. 18, 1895 (**131**).

ALTERNANTHERA, Forsk.

A. straminea (Mart).
Megiphanes straminea, Mart. Island of Cozumel (*Gaumer*, 1885).

GOMPHRAENA, Linn.

G. capitulifiora.
Campo about Izamal, Jan. 23, 1895 (**174**).

G. vermicularis, Linn.
Island of Holbox (*Gaumer*, 1886).

IRISINE, P. Br.

I. celosioides, Linn.
Island of Cozumel (*Gaumer*, 1885 and 1886). Yucatan loc. ignot. (*Johnson*).

I. luzuliflora, Griseb.
Plentiful on the north slope of the Cerro Grande, Izamal, Jan. 13, 1895 (**59**). On the ruins of the Monjas, Chichen Itza, Jan. 17, 1895. In our specimens the stem is ligneous and nearly erect.

PHYTOLACCACEAE.

RIVINA, Linn.

R. humilis, Linn.
Island of Cozumel (*Gaumer*, 1885).

R. humilis glabra, Linn.
Base of smaller mound, Izamal, Jan. 14, 1895 (**85**), in flower and young fruit. Growing in crevices of masonry, old Hacienda, Chichen Itza, Jan. 17, 1895 (**101**), in flower and ripe fruit. Among the rocks, South Point, Island of Mugeres, in flower and ripe fruit Jan. 2, 1895.

R. scandens (Linn.) Mill.
Rivinia humilis scandens, Linn. *Villamilla octandra*, Hook. Yucatan, loc. ignot. (*Johnson*).

R. lævis, Linn.
Islands of Mugeres and Cozumel (*Gaumer*, 1886).

PETIVERIA, Linn.

P. alliacea, Linn.
Yucatan, loc. ignot. (*Linden, Johnson*).

NYCTAGINACEAE.

MIRABILIS, Linn.

M. longifolia, Linn. **XPAK-UM-PAK.**
Frequent about the ruins of Chichen Itza, Jan. 18, 1895 (**121**).

BOERHAAVIA, Linn.

B. erecta, Linn.
Island of Cozumel (*Gaumer*, 1885 and 1886). Campo about Izamal, Jan. 23, 1895 (**184**).

B. hirsuta, Linn.
Campo about Izamal, Jan, 23, 1895 (**172**).

NYMPHACEAE.

CABOMBA, Aubl.

C. aquatica, Aubl.
Yucatan, loc. ignot. (*Linden*).

CASTALIA, Salisb.

C. ampla (DC.) Salisb. **LAB.**
Nymphæa ampla, DC. At the margin of the Aguada of Xcolak, near Izamal, Jan. 22, 1895 (**159**).

ANONACEAE.

ANONA, Linn.

A. squamosa, Linn. *Saramuyo.* Custard Apple. **OALMUY.**
Cultivated extensively in gardens in Merida and Izamal for its fruit.

A. Insularis, Hemsl.
Type locality, Island of Cozumel (*Gaumer*, 1885).

A. muricata, Linn. *Guanabano.*
Cultivated extensively on the Peninsula, Merida, Izamal.

RANUNCULACEAE.

CLEMATIS, Linn.

C. Caripensis, H. B. K.
Clematis Caracasana, DC. Yucatan loc. ignot. (*Johnson*).

C. dioica, Linn.
Edges of the scrub Izamal, Jan. 23, 1895 (**178**), in fruit only. Island of Cozumel (*Gaumer*, 1886).

C. Flammulastrum, Griseb.
Ruins of Uxmal (*Schott*).

LAURACEAE.

PERSEA, Gaertn.

P. gratissima, Gaertn. *Aguacate.* Alligator Pear. **ON.**
A small spreading tree, cultivated extensively for its fruit. Izamal, Merida, Tuncas, Chichen Itza.

NECTANDRA, Roland.

N. Willdenowiana, Nees.
Island of Cozumel (*Gaumer*, 1885), varietates, Hemsley.

CASSYTHA, Linn.

C. filiformis, Linn.
Cassytha Americana, Nees. Island of Holbox (*Gaumer*, 1886).

PAPAVERACEAE.

ARGEMONE, Linn.

A. Mexicana, Linn. *Cardosanto.*
About the huts and streets of Cedral, Island of Cozumel Jan. 5, 1895, (**34**); also collected here by Dr. Gaumer in 1885–6. A garden weed in Merida and Izamal.

CRUCIFERAE.

CAKILE, Gaertn.

C. æqualis, L'Her.
Cakile maritima, Rich. Island of Mugeres, on the sands of the beach together with *Tournefortia gnaphalodes* forming the first verdure back from the sea. North Point and near the Campo Santo of Dolores, Jan. 1, 1895 (**13**), also collected here by Dr. Gaumer in 1885. Progreso, in the same situation and association, Jan. 26, 1895.
A form of this species, Island of Cozumèl (*Gaumer*, 1885).

CAPPARIDACEAE.

CAPPARIS, Linn.

C. amygdalina, Lam.
Yucatan, loc. ignot. (*Linden*).

CRATAEVA, Linn.

C. Tapia, Linn.
Tree 20 to 25 feet; woodlands about Izamal, Jan. 23, 1895 (**187**).

CRASSULACEAE.

BRYOPHYLLUM, Salisb.

B. pinnatum (Lam.) S. Kurz.
Cotyledon pinnatum, Lam. *Bryophyllum calycinum*, Salisb. Plentiful on the north aspect of the Cerro Grande, Izamal, Jan. 13, 1895 (**62**). A dense growth of this species masses the ground about the east margin of the Cenote of Ix-Kil, Chichen Itza. Island of Cozumel (*Gaumer*, 1885).

ROSACEAE.

CHRYSOBALANUS, Linn.

C. Icaco, Linn.
Island of Mugeres (*Gaumer*, 1886).

LEGUMINOSAE.

PITHECOLOBIUM, Mart.

P. ligustrinum, Kl.
Yucatan, loc. ignot. (*Johnson*).

P. sericifolium, Benth. ?
Island of Cozumel (*Gaumer*, 1885), "Vine 10 to 50 feet; common."

P. oblongum, Benth.
Island of Mugeres (*Gaumer*, 1886).

P. Unguis-Cati (Linn.) Benth.
Island of Mugeres (*Gaumer*, 1886). Along the Tepakaam road, near Izamal, Jan. 15, 1895 (**92**).

CALLIANDRA, Benth.

C. Portoricensis (Jacq.) Benth.
Yucatan, loc. ignot. (*Johnson*).

LYSILOMA, Benth. ?

L.———— sp. nov. Hemsl.
Island of Cozumel (*Gaumer*, 1885). Tree 25 to 100 feet; abundant, fruit wanting. Aff. *L. Schiedeanae*, Hemsley.

ACACIA, Linn.

A. cornigera (Linn.) Willd. ?
Mimosa cornigera, Linn. Very abundant in the scrub lands of the Island of Cozumel, and from Progreso to Chichen Itza.

A. Farnesiana (Linn.) Willd. ?
Mimosa Farnesiana, Linn. Island of Cozumel (*Gaumer*, 1885 and 1886). "Not common."

A. spadicigera, Ch. & Schl.
Island of Cozumel (*Gaumer*, 1885). "Abundant." Dr. Gaumer tells me that the Cozumelian plant is the same as that of the Main, and that his plants from there differ in no way from my No. **221** from Izamal. This removes the question mark from the species in Biol. Centr. Am., Vol. 4, p. 99.

LEUCAENA, Benth.

L. glauca (Linn.) Benth.
Mimosa glauca, Linn. Campo about Izamal, Jan. 23, 1895 (**182**).

MIMOSA, Linn.

M. trijuga, Benth.
Yucatan, loc. ignot. (*Johnson*).

DESMANTHUS, Willd.

D. depressus, Humb. & Bonpl.
Island of Cozumel (*Gaumer*, 1886).

TAMARINDUS, Linn.

T. Indica, Linn.
Cultivated at Cedral, Island of Cozumel.

BAUHINIA, Linn.

B. divaricata, Linn. *Pata de Vaca.*
Bauhinia porrecta, Sw. Scrub about Izamal, Jan. 22, 1895 (**216**). Yucatan, loc. ignot. (*Johnson*). Island of Cozumel (*Gaumer*, 1885 and 1886).
The Spanish name "*Pata de Vaca*" is applied to all of the Bauhinias having bilobed leaves.

B. latifolia, Cav.
Scrub about Izamal, Jan. 22, 1895 (**217**).

B. Lamarckiana, DC.
Borders of the more open woods, Mascab Pixoy, Jan. 22, 1895 (**144**).

CASSIA, Linn.

C. biflora, Linn.
Izamal, Jan. 23, 1895 (**186**), Bushy, 4 ft. high.

C. alata, Linn.
Island of Cozumel (*Gaumer*, 1885), "Rare."

C. emarginata, Linn.
Island of Cozumel (*Gaumer*, 1885), "very abundant."

C. ——— sp. **YAXHABIN.**
Tree 10 to 20 feet; abundant in the scrub-land from Progreso to Merida along the railroad. Flowers bright yellow, the most striking feature of the flora of January.

C. bacillaris, Linn. f.
Yucatan, loc. ignot. (*Johnson*).

C. bicapsularis, Linn.
Yucatan, loc. ignot. (*Johnson*).

C. reticulata, Willd.
Yucatan, loc. ignot. (*Johnson*).

C. occidentalis, Linn.
Lagoona Colombia, Island of Cozumel; Izamal, and Chichen Itza, Jan. 1895. Island of Mugeres (*Gaumer*, 1886).

C. ——— sp. (Hemsl).
Islands of Cozumel and Mugeres (*Gaumer*, 1886).

HAEMATOXYLON, Linn.

H. Campechianum, Linn.
Yucatan, loc. ignot. (*Johnson*).

CAESALPINIA, Linn.

C. Bonducella (Linn). Flem.
Guilandina Bonducella, Linn. A mass of this species on the skirts of the woods, South Point, Island of Mugeres, in fruit only Jan. 2, 1895.

C. pulcherrima (Linn). Sw.
Poinciana pulcherrima, Linn. Island of Cozumel (*Gaumer*, 1885), "very abundant."

C. exostemma, DC.
In the scrub along the road to Chal-tun-ha near Izamal, Jan. 13, 1895 (**75**).

ATELEIA, Moc. & Sess.

A. pterocarpa (DC.)
Edge of woods, road to Cedral, Island of Cozumel, Jan. 5, 1895 (**44**).

A. microcarpa (Pers).
Pterocarpus microcarpus, Pers. Scrub about Izamal, Jan. 23, 1895 (**177**).

SOPHORA, Linn.

S. tomentosa, Linn.
Island of Mugeres (*Gaumer*, 1886).

CROTALARIA, Linn.

C. pumila, Orteg.
Island of Mugeres (*Gaumer*, 1886).

INDIGOFERA, Linn.

I. Anil, Linn.
Island of Cozumel (*Gaumer*, 1886).

I. mucronata, Spreng.
Island of Cozumel (*Gaumer*, 1886).

PAROSELA, Cav.

P. Domingensis (DC).
Dalea Domingensis, *DC.* In the scrub bordering the road to Chal-tun-ha, near Izamal, Jan. 13, 1895 (**77**).

CRACCA, Linn.

C. cinerea (Linn), O.K.
Galega cinerea, Linn. *Tephrosia cinerea*, Pers. Yucatan, loc. ignot. (*Linden*). Island of Cozumel (*Gaumer*, 1885).

C. Caribæa, Benth.?
Island of Cozumel (*Gaumer*, 1885).

C. ———— sp. (Hemsl.)
Island of Cozumel (*Gaumer*, 1885), "a fragment only."

GLIRICIDIA, H.B.K.

G. ———— sp. nov. Hemsl.
Island of Cozumel (*Gaumer*, 1885). "Tree 50 feet; not abundant. In the absence of the fruit the genus of this tree cannot be satisfactorily determined." Hemsley.

G. maculata, H.B.K.
Island of Cozumel (*Gaumer*, 1886).

DIPHYSA, Jacq.

D. ———— sp. nov.? Hemsl.
Island of Cozumel (*Gaumer*, 1885), "Specimen insufficient for satisfactory determination. It may be: *D. Sennoidei* var. nov.? Hemsley."

AESCHYNOMENE, Linn.

A. Americana, Linn.
Yucatan, loc. ignot. (*Linden*).

MEIBOMIA, Adans.

M. affinis (Schl.) O.K.
Desmodium affine, Schl. Yucatan, loc. ignot. (*Johnson*).

M. adsendens (Sw.) O.K.
Hedysarum adsendens, Sw. Island of Cozumel (*Gaumer*, 1885).

AMERIMNON, P. Br.

A. Campechianum (Benth) O.K.
Dalbergia Campechiana, Benth. Yucatan, loc. ignot. (*Linden, Johnson.*)

LONCHOCARPUS, H.B.K.

L. violaceus, H.B.K.?
Island of Cozumel (*Gaumer*, 1885). "Tree 20 to 60 feet; very abundant."

ICHTHYOMETHIA, P. Br.

I. Piscipula (Linn.) O.K.
Erythrina Piscipula, Linn., *Piscidia Erythrina*, Linn. Island of Cozumel (*Gaumer*, 1885), "very common."

BRADBURYA, Raf

B. Virginiana (Linn.) O.K.
Clitoria Virginiana, Linn., *Centrosema Virginianum*, Benth. Island of Cozumel (*Gaumer*, 1885).

B. pubescens (Benth.) O.K.
Centrosema pubescens, Benth. Yucatan, loc. ignot. (*Johnson*).

ERYTHRINA, Linn.

E. carnea, Ait.
Frequent in the scrub about Izamal, Jan. 14, 1895 (**94**).

CALOPOGONIUM, Desv.

C. caeruleum, Desv.
Yucatan, loc. ignot. (*Johnson*).

GALACTIA, P. Br.

G. ——— sp. (Hemsl.)
Island of Cozumel (*Gaumer*, 1886).

RHYNCHOSIA, Lour.

R. minima (Linn.) DC.
Dolichos minima, Linn. Island of Mugeres (*Gaumer*, 1886).

PHASEOLUS, Linn.

P. lunatus, Linn.
Island of Cozumel (*Gaumer*, 1886).

P. vulgaris, Linn. *Frijol.* **BUUL.**
Island of Cozumel (*Gaumer*, 1886).

P. Truxillensis, H.B.K.
Yucatan, loc. ignot. (*Johnson*).

DOLICHOS, Linn.

D. puriens, Linn. **CHIICAN.**
In the thicket, base of the smaller mound, Izamal, Jan. 14, 1895, in fruit only.

MALPIGHIACEAE.

HIRAEA, Jacq.

H. Barclayana, Benth. ?
Island of Cozumel (*Gaumer*, 1885).

HETEROPTERYS, Kth.

H. floribunda, H.B.K.
Yucatan, loc. ignot. (*Johnson*).

H. Lindeniana, A. Juss.
Yucatan, typ. loc. ignot. (*Linden*).

H. longifolia, H.B.K.
Yucatan, loc. ignot. (*Johnson*).

MALPIGHIA, Linn.

M. undulata, Juss.
Island of Cozumel (*Gaumer*, 1886).

BUNCHOSIA, Rich.

B. Lindeniana, A. Juss.
Yucatan, loc. ignot. (*Johnson*).

BYRSONIMA, Rich.

B. Karwinskiana, A. Juss.
Yucatan, loc. ignot. (*Johnson*).

B. crassifolia, H.B.K., var. (Hemsl.)
B. Cumingiana, A. Juss. Island of Cozumel (*Gaumer*, 1885). "Not common."

ZYGOPHYLLACEAE.

TRIBULUS, Linn.

T. terrestris cistoides (Linn.) Oliver. *Abrojo.* **CHANXNUC.**
Tribulus cistoides, Linn. Streets of Progreso, May, 1887, Jan. 26, 1895 (**200**).

T. maximus, Linn.
Island of Cozumel (*Gaumer*, 1885).

RUTACEAE.

RUTACEARUM, Hemsl.

R. ——— Gen. et. sp. nov. Hemsl.
Island of Mugeres (*Gaumer*, 1886). "This may prove the type of a new genus; but without fruit it is impossible to be sure on this point." Hemsley.

ZANTHOXYLUM, Linn.

Z. pterota, Linn.
Island of Cozumel (*Gaumer*, 1885). "Not common."

ENSENBECKIA, Kth

E. pentaphylla (Macf.) Griseb.
Island of Mugeres (*Gaumer*, 1886).

PTELEA, Linn.

P. pentandra, DC.

A low tree in the shrubbery of the north side of the smaller mound at Izamal, Jan. 14, 1895 (**86**). Immature.

CHALCAS, Linn.

C. exotica (Linn).

Murraya exotica, Linn. Introduced on the Island of Cozume (*Gaumer,* 1885).

CITRUS, Linn.

C. aurantium, L. *Naranja.* Orange.

Cultivated throughout the mainland, and on the Island of Cozumel. The first planting of the orange in Mexico was doubtless that of Bernal Diaz, who, while attached to the army of Cortez, sowed seed near the present location of Vera Cruz, and transplanted the resulting trees after the conquest. Several of these trees were again transplanted to Merida by Montejo, early in the 16th century.

C. vulgaris, Risso. *Cajera.* Bitter Orange. **SUTZ-PAKAL.**

Cultivated at San Miguel, Island of Cozumel, and on the mainland. Notwithstanding its bitter acid taste it is a very refreshing fruit, and is much prized by the natives.

C. Limonum, Risso. *Limon.* Lemon.

Cultivated at Progreso, Merida and Izamal, not so extensively however as the next species.

C. Limetta, Risso. *Limoncillo.* Lime.

Cultivated throughout the mainland, and on the Island of Cozumel; met with as far south as Mascab Pixoy.

SIMARUBACEAE.

SURIANA, Linn.

S. maritima, Linn.

North Point, Island of Mugeres, Jan. 2, 1895 (**23**). A shrub 5 to 10 feet high, bark thin, grayish-white, heart wood red, very hard. A prominent feature of the littoral growth.

Islands of Holbox and Mugeres (*Gaumer,* 1886).

PICRAMNIA, Sw.

P. Antidesma, Sw.

Island of Cozumel (*Gaumer,* 1885).

BURSERACEAE.

BURSERA, Linn.

B. Simaruba (Linn.) Sarg. **CHACAH.**

Bursera gummifera, Jacq. Abundant on the Islands of Mugeres and especially Cozumel, also between Tuncas and Chichen Itza on the Main, where its peculiar coppery trunk and divaricate branches render it a conspicuous object in the forest. It grows from 25 to 50 feet. Not yet in leaf Jan. 1-22, 1895.

MELIACEAE.

MELIA, Linn.

M. Azederach, Linn.
Introduced on the Island of Cozumel (*Gaumer*, 1885).

TRICHILIA, P. Br.

T. Terminalis, Jacq.
Island of Cozumel (*Gaumer*, 1885). "Not common."

CEDRELA, P. Br.

C. odorata, Linn. **KUCHÉ. KULCHÉ.**
The *Cedro* or "Cedar" of Mexico. Common in the interior at Chichen Itza. Often growing from 20 to 40 feet high.

C. Glaziovii, C. DC.
In the interior forests (*Linden*).

POLYGALACEAE.

POLYGALA, Linn.

P. paniculata, Linn.
Yucatan, loc. ignot. (*Johnson*).

EUPHORBIACEAE.

PHYLLANTHUS, Linn.

P. acuminatus, Vahl.
Yucatan, loc. ignot. (*Linden*).

P. ——— sp. (Hemsl.)
Island of Cozumel (*Gaumer*, 1885). "May be a variety of *P. nutans*." Hemsley.

P. Niruri, Linn.
Island of Cozumel (*Gaumer*, 1886).

CROTON, Linn.

C. maritimus, Walt.
Trailing extensively in the clear sands of the beach at Progreso, Jan. 26, 1895 (**201**). Island of Holbox (*Gaumer*, 1886).

C. ——— sp. (Hemsl).
Island of Cozumel (*Gaumer*, 1885). "May be a variety of *C. astroites*." Hemsley.

C. Zalapensis, H.B.K.
Along the Tuncas road, near Mascab Pixoy, Jan. 22, 1895 (**150**).

ACALYPHA, Linn.

A. setosa, A. Rich.
A low form. Campo about Izamal, Jan. 23, 1895 (**180**).

A. alopocuroides, Jacq.
Island of Cozumel (*Gaumer*, 1886).

RICINUS, Linn.

R. communis, Linn. Higuerilla. Castor Bean. **XOX.**
Very common about dwellings and in waste grounds where it has apparently escaped. Noted at all points on our route on the Peninsula. Fine large plants at Ouan, Jan. 24, 1895 (**194**), Merida, April 14, 1887. Islands of Mugeres and Cozumel.

JATROPHA, Linn.

J. aconitifolia, Mill. *Picar.*
Frequent throughout the dry scrub about Izamal, Jan. 13, 1895 (**66**). On the Island of Cozumel, and at all points on the Peninsula, between Progreso and Chichen Itza.
The Spanish name *Picar*, as applied to this plant, is more a word of warning than a name.

EUPHORBIA, Linn.

(*Anisophyllum-Hypericifoliæ*).

E. pilulifera, Linn.
One small specimen only, collected on the summit of the Monjas, Chichen Itza, Jan. 17, 1895 (**219**). Collected by Dr. Gaumer on the Island of Cozumel, 1885.

E. hypericifolia, Linn.
The common West Indian form, Merida, Jan. 24, 1895 (**197-9**). Streets of San Miguel and Cedral, Island of Cozumel, Jan. 5, 1895. Also collected on Cozumel by Dr. Gaumer, 1885 and 1886.

(*Anisophyllum-Chamaesyceæ*).

E. ——— sp.
An interesting erect or prostrate form, near E. serrula and doubtless new; the specimens too young for determination.
Grassy edge of sands of West beach, Island of Cozumel at the portage to the Lagoon of Colombia, Jan. 7, 1895 (**49, 53**).

E. ovalifolia, Engelm.
A form with less turbinate involucres having no hairs in the throat, otherwise agreeing with the species as described.
Sandy roadside, Dolores, Island of Mugeres, Jan. 1, 1895 (**19**).

E. maculata, Linn.
A very pubescent form, without the usual maculate leaves common on the south point of Mugeres Island, and on the lower courses of "The Watch Tower" ruin, Jan. 2, 1895 (**24**).

E. adenoptera, Bertol.

Fine specimens with large rose-colored appendages on the dryest limestone ledges of the campo, Chichen Itza, Jan. 17, 1895 (**107**).

(*Anisophyllum-Sclerophyllæ*).

E. buxifolia, Lam.

Great tufts of this species form the only break in the wide sandy beach of the east shores of the Island of Cancun. Very common, Jan. 3, 1895 (**31**). Plants 15 to 18 cm. high. Grassy sands of the west shore Island of Cozumel, Jan. 7, 1895 (**52**). Plants 5 to 10 dm. high and very shrubby. Sands of shore at Progreso, Jan. 26, 1895 (**202**). Sandy streets village of Dolores, Island of Mugeres, Jan. 2, 1895 (**20**).

This species was also collected on Mugeres and Holbox Islands by Dr. Gaumer in 1886.

(*Adenopetalum-Cyttarospermum*).

E. Armourii. sp. nov. (Plate II.)

Caule erecto glabro, ramis tenuibus frutescentibus (?), ramulis axillaribus solitariis folio longioribus, partes junioribus et nervos foliorum minute hirtula, foliis longe petiolatis e basi cuneata ellipticis integris acutis anguste lanceolatis, involucris in axilis super. terminalibusque longiuscule pedicillatis breviter turbinatis, lobis truncatis profunde 4-dentatis, glandulis 5 planis transverse ovatis, appendice alba bipartita laciniis oblongo-linearibus obtusis glandula duplo longioribus, stylis profunde bipartitis, capsula glabra, seminibus ovatis ad margines puncto-fevearum prominente tuberculatis. Folia 14-24 mm. lon., 10-19 mm. lat., caules 2½-4 dm. lon., 1-2 mm. diam., invol. 2 mm. lon., 1 mm. lat., semin. 1 mm. lon. ¾ mm. lat.

Growing in the crevices of the masonry forming one of the temples of the sacred Itzae, Chichen Itza, Jan. 17, 1895 (**108**).

Named in honor of Mr. Allison V. Armour, the projector of the expedition on which this collection was made.

E. astroites heterappendiculata, var. nov.

Caule tenui *flacciduli* alternatim ramosissimo, ramis glanduloso-hirtis pluries et divaricatim dichotomis, foliis oppositis minimis *longe petiolatis* orbiculatis vel transverse latioribus adpressiuscule hirtis, stipulis obsoletis, involucris terminalibus pedicillatis valde glanduloso-hirtis, *lobis truncatis* 4-*dentatis*, glandulis transverse ovatis concavis, *appendice integris* 1–4 *dentatis*, *dentato*-2-3 *partitis vel* 4-*partitis*, *laciniis lineari-lanceolatis obtusis*, capsula hirta, stylis profunde bipartitts, seminibus ovatis ad margine puncto-fovearum prominente tuberculatis.

Caules 2–3 dm. lon., ramis 8–12 cm. lon., foliis inferioris 6 mm. lat., invol. 1 mm. lon., semin. 1 mm. lon. ¾ mm. lat. The species, *E. astroites*, F & M., was collected at Tampocoala, Mexico, by Karwinsky.

Crevices of rocks, east slope of the Cerro Grande, Izamal, Jan. 13, 1895 (**64**).

EUPHORBIA ARMOURII, sp. nov.

(*Adenopetalum-Alectoroctonum*).

E. petiolaris, Sins.
Collected on the Island of Cozumel by Dr. Gaumer in 1886.

(*Poinsettia*).

E. heterophylla, Linn. **HOBONKAK**
The usual form of the species. The Maya name "Hobonkak" signifies "Fire in the hollow" referring to the orange-red bases of the floral leaves.
Island of Mugeres, Jan. 1, 1895; Island of Cozumel, Jan. 5, 1895; Chichen Itza, Jan. 17, 1895 (**111**); Progreso, Jan. 26, 1895. Collected on Cozumel by Dr. Gaumer in 1885.

(*Tithymalus-Ipecacuanhæ*)

E. trichotoma, Kunth.
Collected on the Island of Cozumel by Dr. Gaumer in 1885.

MALVACEAE.

ABUTILON, Gaertn.

A. Indicum (Linn.) Don. **SAC-XIEU.**
Sida Indica, Linn. Frequent on the campo about the ruins, Chichen Itza, Jan. 18, 1895 (**124**).

MALVEOPSIS, Presl.

M. spicata (Linn.) O.K. **CHÉ-CHÉ-BÉ.**
Malva spicata, Linn. *Malvastrum spicatum*, Griseb. Campo about the ruins, Chichen Itza, Jan. 18, 1895 (**124**).
The Maya name Che-che-be, signifying "The Home of the Garrapata," is commonly applied to all malvaceæ resembling this species.

SIDA, Linn.

S. acuta carpinifolia (Linn. f.) K. Schum.
Island of Cozumel (*Gaumer*, 1885).

ANODA, Cav.

A. hastata (Willd.) Cav. *Amapolite.*
Sida hastata, Willd. About the ruins of the old church of the Hacienda, Chichen Itza, Jan, 18, 1895 (**126**).

A. acerifolia (Zucc.) DC.
Sida acerifolia, Zucc. About the base of the smaller mound, Izamal, Jan. 14, 1895 (**83**).

MALVAVISCUS, Adans.

M. sepium, Schl.
Yucatan, loc. ignot. (*Johnson*).

M. ——— sp. (Hemsl.)
Island of Cozumel (*Gaumer*, 1886). "Near *M. pilosus*, DC." Hemsley.

M. arboreus, Cav. forma.
A common shrub about Izamal. On the Cerro Grande, Jan. 13, 1895, (**60**). Fruits of this species were collected afterwards at Silam by Dr. Gaumer (April, 1895).

M. concinnus, H.B.K.
Roadside and edges of open woods. Cedral, Island of Cozumel, Jan. 5, 1895 (**42**).

HIBISCUS, Linn.

M. tubiflorus, DC. **HOL.**
Shrub 4 to 6 feet; widely branching. Road from Mascab Pixoy to Xuku, in flower and fruit Jan. 22, 1895 (**157**).

BOMBACACEAE.

BOMBAX, Linn.

B. Ceiba, Linn. *Ceiba,* **YAXCHÉ.**
One of the largest trees noted on the trip, 80 to 100 feet. Fine specimens in and about Izamal, and between Xcolak and Tuncas, also at Chichen Itza.

CAROLINEA, Linn.

C. alba (Aubl.) Lodd. **XCUYCHÉ. KUYCHÉ.**
The beautiful *Amapola blanca,* cultivated extensively about Izamal and Merida. Not yet in bloom Jan. 21, 1895.

C. fastuosa, (D.C.) Sesse. *Amapola colorado.*
Cultivated with the last. A few of the magnificent rose-colored flowers expanded Jan. 21, 1895.

C. macrocarpa (Aubl.) Ch. & Schl.
Pachira macrocarpa, Aubl. Yucatan, loc. ignot. (*Johnson*).

CEIBA, Medic.

C. pentandra, (Linn.) Gaertn. *Pochote,* **PEEM.**
Bombax pentandrum, Linn. *Bombax Ceiba,* Lun. (non Linn.) *Eriodendron anfractuosum,* D.C. Plentiful throughout the Peninsula. Fine specimens both to the north and south of Izamal, also at Mascab Pixoy.

STERCULIACEAE.

MELOCHIA, Linn.

M. pyramidata, Linn.
Island of Cozumel (*Gaumer*, 1885).

M. tomentosa, Linn. ?
Island of Cozumel (*Gaumer*, 1885). "This may be a distinct species; it is the same as Berlandier's 3032 from the Texano-Mexican region, and there is also a specimen in the Kew Herbarium, collected by Coulter, from Sonora Alta." Hemsley.

WALTHERIA, Linn.

W. Americana, Linn.
Island of Cozumel ? (*Gaumer*, 1886). In a small clearing on the Tuncas road near Mascab Pixoy, Jan. 22, 1895 (**152**). Along the railroad at Acalchen, Jan. 24, 1895 (**192**).

AYENIA, Linn.

A. magna, Linn.
Yucatan, loc. ignot. (*Linden*).

THEOBROMA, Linn.

T. Cacao, Linn. *Cacao.*
A small tree, cultivated in favored localities for its fruit, the chocolate-nut. Chichen Itza, in the Cenote de Cafetal.

GUAZUMA, Plum.

G. tomentosa, H.B.K. **PIXOY.**
On the road to Cedral, Island of Cozumel, in fruit, Jan. 5, 1895 (**40**). At Mascab Pixoy, Luché Pixoy, and Chichen Itza, Island of Cozumel (*Gaumer*, 1885). Yucatan, loc. ignot. (*Johnson*).

STERCULIA, Linn.

S. Carthagensis, Cav.
Yucatan, loc. ignot. (*Johnson*).

GUTTIFERAE.

MAMMEA, Linn.

M. Americana, Linn.
Island of Cozumel (*Gaumer*, 1885). "Rare."

BIXACEAE.

MAXMILIANEA, Mart. & Schr.

M. hibiscoides (Humb. & Bonpl.) **CHUM.**
Cochlospermum hibiscoides, Humb. & Bonpl. A tree 30 to 40 feet high, with a thick bark and very soft wood. The bark is used in the manufacture of "Cubos"; a sort of bucket. At Xuku, Jan. 22, 1895 (**156**). In flower, not in leaf.

MYROXYLON, Forst.

M. nitidum (Schl.)
Hisingera nitida and *pubescens*, Schl. *Xylosma nitidum.* A. Gray. Yucatan, loc. ignot. (*Johnson*).

BIXIA, Linn.

B. Orellana, Linn. *Arnatto.* **URUCU.**
Cultivated for its seed, the aril of which yields the "Butter Color" of Commerce. Merida, Izamal, in fruit along the road to Chal-tun-ha, Jan. 14, 1895 (**81**).

VIOLACEAE.

CALYPTRION, Ging.

C. Hybanthus (Aubl.)
Viola Hybanthus Aubl. *Corynostylis Hybanthus* (Aubl.) Mart. & Zucc., *Calyptrion Aubletii* (E. Mey.) Ging. Yucatan, loc. ignot. (*Johnson*).

ANACARDIACEAE.

RHUS, Linn.

R. Metopium, Linn.
Island of Cozumel (*Gaumer*, 1885). "Rare."

SPONDIAS, Linn.

S. lutea, Linn. *Abal.* **XKININ-HOBÓ.**
Cultivated on the Island of Cozumel and throughout the Peninsula. Yucatan, loc. ignot. (*Johnson*). The fruit is known as the *Ciruela* or Yucatan Plum.

S. ——— sp. (Hemsl.)
Island of Cozumel (*Gaumer*, 1885). "Only male flowers and quite young leaves." (Hemsley).

MANGIFERA, Linn.

M. Indica, Linn. *Mango.*
Island of Cozumel (*Gaumer*, 1886).

CELASTRACEAE.

CASSINE, Linn.

C. ——— sp. (Hemsl.)
Elaeodendron sp. Hemsl. Island of Mugeres (*Gaumer*, 1886).

CROSSOPETALUM, P. Br.

C. pallens (Smith).
Myginda pallens, Smith. Island of Cozumel (*Gaumer*, 1885). "Very rare."

HIPPOCRATEACEAE.

HIPPOCRATEA, Linn.

H. tenella, Miers. ?
Island of Cozumel (*Gaumer*, 1885). "This may be specifically different from the Mexican plant, but the material is insufficient to settle the question." Hemsley.

SAPINDACEAE.

SERIANA, Schum.

S. Mexicana (Linn.) Willd.
Paulinia Mexicana, Linn., *Serjania Mexicana*, Willd. Yucatan, loc. ignot. (*Johnson*).

S. plicata (Radlk).
Yucatan, loc. ignot. (*Johnson*).

S. scatans (Radlk).
Yucatan, loc. ignot. (*Johnson*).

SCHMIDELIA, Linn.

S. comina, Sw. ?
Yucatan, loc. ignot. (*Linden*).

PAULINIA, Linn.

P. curassavica, Linn.
Island of Cozumel (*Gaumer*, 1886).

SAPINDUS, Linn.

S. ——— sp.
At the Lagoona Colombia, Island of Cozumel, Jan. 7, 1895, in fruit only. Also collected here by Dr. Gaumer in 1886.

ALVARADOA, Liebm.

A. amorphoides, Liebm. **BEL-CINICHÉ.**
In the scrub bordering the Chal-tun-ha road, near Iżamal, Jan. 13, 1895 (**76**).

RHAMNACEAE.

GOUANIA, Jacq.

G. stipularis, DC.
In the interior forests (*Linden*); type locality?

VITACEAE.

VITIS, Linn.

V. sicyoides monstrosa, Baker.
Yucatan, loc. ignot. (*Johnson*).

V. tiliaefolia, Willd.
Island of Cozumel (*Gaumer*, 1885).

TILIACEAE.

TRIUMFETTA, Linn.

T. althaeoides, Lam.
Yucatan, loc. ignot. (*Johnson*)

CORCHORUS, Linn.

C. siliquosus, Linn.
Yucatan, loc. ignot. (*Johnson*).

PASSIFLORACEAE.

PASSIFLORA, Linn.

P. Andersonii, DC. ?
Island of Cozumel (*Gaumer*, 1885) "Vine 30 to 50 feet; very abundant."

P. foetida, Linn.
Island of Cozumel (*Gaumer*, 1885), Island of Mugeres (*Gaumer*, 1886).

CARICACEAE.

CARICA, Linn.

C. Papaya, Linn. (Plate III.) *Papaya los Pajaros.* **CHICH PUT.**
Frequent on the islands of Mugeres, Cankun, and Cozumel, in fruit, very few specimens in flower. Chichen Itza, Jan. 20, 1895 (**143**). Island of Cozumel (*Gaumer*, 1886).
This is the native and natural form, called *Papaya los Pajaros* or "Bird Papaya" by the Spaniards, and Chich Put or "Little Papaya" by the Mayas. The fruit of this form is small, about one inch in diameter, nearly globular and not edible.

C. Papaya, Linn. (Cultivated) (Plate IV.) *Papaya.* **PUT.**
Commonly cultivated throughout the Peninsula, and on the Island of Cozumel, producing large pear-shaped fruit from 12 to 16 inches long and 6 to 9 inches in diameter at the larger end. The pulp is of an orange-salmon color, rich, juicy and delicious. Izamal, Jan. 13, 1895 (**69**).
The flavor of the fruit here, is said to be largely dependent upon the quantity of seed in the individual, those having but few seeds being far finer in every feature than those having many. It is also said that in raising Papayas from the seed in this locality selection plays no part, as the result can not be known until the plant bears its fruit; some of the seeds will revert to the original of the species (*Papaya los Pajaros*) while others will produce excellent Papayas.

CARICA PAPAYA, Linn.
(*Natural.*)

CARICA PAPAYA, Linn.
(*Cultivated.*)

JACARATIA, Marcgr.

J. Mexicana, DC. *Bonete,* **KUMCHÉ.**

A large tree 15 to 40 feet high, abundant between Merida and Izamal. A few individuals in flower at Izamal, Jan. 15, 1895 (**96.**)

The somewhat triangular fruit, called Bonete, has a yellow mawkish pulp, which is usually eaten in a sort of custard.

LOASACEAE.

GRONOVIA, Linn.

G. scandens, Linn. **LAL-MUCH.**

Extensively trailing over the campo about the ruins of Chichen Itza, Jan. 18, 1895 (**128**).

CACTACEAE.

CEREUS, Haw.

C. flagelliformis (Linn.) Mill. **CANCHOH.**

Island of Mugeres on the south-western coast rocks. Between Tuncas and Mascab Pixoy, and at Chichen Itza.

C. grandiflorus (Linn.) Haw. *Pitaya.*

Plentiful on the Islands of Mugeres and Cozumel. Between Tuncas and Chichen Itza.

C. ——— sp. (Hemsley).

On trees, Island of Cozumel (*Gaumer,* 1885). An *C. nycticalus* or *C. MacDonaldiæ*? Hemsley.

OPUNTIA, Adans.

O. Tuna (Linn.) Mill. *Tuna, Nopal.* **PAKAN.**

Cactus Opuntia, Linn. A large patch on the south point of the Island of Mugeres, bearing the largest and finest fruits that I have ever seen. In fruit Jan. 2, 1895. Frequent on the mainland, especially between Progreso and Merida.

LYTHRACEAE.

LAWSONIA, Linn.

L. alba, Lam.

Yucatan loc. ignot. (*Johnson*). Island of Cozumel (*Gaumer,* 1885). Of this species Hemsley says: "Introduced from the Old World."

RHIZOPHORACEAE.

RHIZOPHORA, Linn.

R. Mangle, Linn. **TAPCHÉ.**

Plentiful about the Lagoona Colombia, Island of Cozumel, where it forms many small "islands." Plentiful at the northwest end of the Island of Cankun, about which are also many "islands" composed entirely of this species. Island of Holbox (*Gaumer,* 1886).

MYRTACEAE.

PSIDIUM, Linn.

P. Guajava, Linn. *Guayabo*, **PICHI.**
Island of Cozumel (*Gaumer*, 1885). Cultivated at Merida, Izamal, and Tuncas.

PIMENTA, Lindl.

P. vulgaris, Wight.
Myrtus Pimenta, Linn. Beautiful large specimens in cultivation in the Cenote de Cafetal, Chichen Itza.

EUGENIA, Linn.

E. ——— sp. nov.? Hemsl.
Island of Cozumel (*Gaumer*, 1885). "Quite different from anything in the Kew Herbarium." Hemsley.

COMBRETACEAE.

BUCERAS, P. Br.

B. Catappa (Linn.) Hitch.
Terminalia Catappa, Linn. Island of Cozumel (*Gaumer*, 1885).

CONOCARPUS, Linn.

C. erecta, Linn.
Island of Cozumel (*Gaumer*, 1885). Island of Holbox (*Gaumer*, 1886).

MELASTOMACEAE.

TIBOUCHINA, Aubl.

T. longifolia (Vahl.)
Rhexia longifolia, Vahl. *Pleroma longifolia*, Hemsl. Yucatan loc. ignot. (*Johnson*).

CONOSTEGIA, G. Don.

C. Xalapensis (Bonpl.) G. Don.
Melastoma Xalapensis, Bonpl. Yucatan, loc. ignot. (*Johnson*).

CLIDEMIA, G. Don.

C. petiolaris, Schl.
Yucatan, loc. ignot. (*Johnson*).

MYRSINACEAE.

JACQUINIA, Linn.

J. Caracasana, H.B.K.
Island of Cozumel (*Gaumer*, 1885).

J. aristata, Jacq.
Island of Cozumel (*Gaumer*, 1885). Island of Holbox (*Gaumer*, 1886).

PARATHESIS, Hook f.

P. crenulata, Hook f.
Yucatan, loc. ignot. (*Johnson*).

PLUMBAGINEAE.

PLUMBAGO, Linn.

P. scandens, Linn.
Base of smaller mound, Izamal, Jan. 14, 1895 (**84**).

SAPOTACEAE.

ACHRAS, Linn.

A. Sapota, Linn. *Sapote.* **TAUCH. YA.**
Widely cultivated for its delicious fruit, which also yield, on puncture, the finest of the Yucatan Gum or "*Chicle.*" Also natural to many parts of the Peninsula, particularly in the eastern section where it is abundant and often reaches a growth of 50 to 100 feet in height.
Merida, Izamal, Yucatan loc. ignot. (*Linden*).

BUMELIA, Sw.

B. retusa, Sw. var? (Hemsl.)
Island of Cozumel (*Gaumer*, 1885). Island of Mugeres (*Gaumer*, 1886).

LUCUMA, Molina.

L. mammosa, Gaertn. **CHACHAL HAAS.**
A large and beautiful tree, cultivated for its fine fruit the *Mamey Colorado.* The large seed has a powerful amygdalous odor.
Izamal, Merida, Tuncas, Chichen Itza.

GENTIANACEAE.

EUSTOMA, Don.

E. exaltatum (Linn). Griseb.
Gentiana exaltata, Linn. Island of Cozumel (*Gaumer*, 1885). Island of Holbox (*Gaumer*, 1886).

APOCYNACEAE.

VALLESIA, R. & Pav.

V. glabra, Cav.
Island of Mugeres (*Gaumer*, 1886).

RAUWOLFIA, Linn.

R. heterophylla, Roem & Sch.
Yucatan loc. ignot. (*Johnson*).

THEVETIA, Linn.

T. nitida, A. DC.
Yucatan loc. ignot. (*Johnson*).

T. neriifolia, Juss.
Island of Cozumel (*Gaumer*, 1885 and 1886).

T. Gaumeri, Hemsl.
Type locality, Island of Cozumel (*Gaumer*, 1885), "Tree 40 to 60 feet high; not common."

PLUMERIA, Linn.

P. alba, Linn. **SABACNICTÉ.**
Suburbs of Merida, frequent May, 1887.

TABERNAEMONTANA, Linn.

T. amygdalifolia, Jacq. **UOUPEK.**
Along the road to Valladolid beyond Tuncas. Not in flower, Jan. 17, 1895.

T. Acapulcensis, Miers.
Island of Cozumel (*Gaumer*, 1885).

ASCLEPIADACEAE.

PHILIBERTIA, H. B. K.

P. Lindeniana (Decne). Hemsl.
Sarcostemma Lindenianum, Decne. Yucatan type loc. ignot. (*Linden*).

MACROSCEPIS, H. B. K.

M. rotata, Decne.
Yucatan, type loc. ignot. (*Linden*).

ASCLEPIAS, Linn.

A. curassavica, Linn. **CUCHILXIU.**
Along the railroad, at the station of Ouan, Jan. 24, 1895 (**195**), fruits only. Island of Cozumel (*Gaumer*, 1885).

A. pratensis, Benth.
Along the railroad, at the station of Acalchen, Jan. 24, 1895 (**193**). Differs from the description in having roughly pubescent leaves, especially on the veins above and beneath.

METASTELMA, R. Br.

M. parviflorum (Sw.) R. Br.
Cynanchum parviflorum, Sw. Islands of Mugeres and Holbox (*Gaumer*, 1886).

ENSLENIA, Nutt.

E. racemosa (Jacq)
Cynanchum racemosum, Jacq. *Roulinia Jacquini*, Decne. Yucatan, loc. ignot. (*Linden*).

MARSDENIA, R. Br.

M. maculata, Hook.
Island of Cozumel (*Gaumer*, 1885).

CONVOLVULACEAE.

EVOLVULUS, Linn.

E. nummularius (Linn.) Linn.
Yucatan, loc. ignot. (*Johnson*).

IPOMOEA, Linn.

I. Pes-caprae (Linn.) Sweet.
Convolvulus Pes-caprae, Linn. Island of Cozumel (*Gaumer*, 1885). "Sometimes attains an hundred feet in length."

I. triloba, Linn.
Island of Cozumel (*Gaumer*, 1886).

I. sidaefolia, Linn.
Island of Cozumel (*Gaumer*, 1886).

I. coccinea, Linn.
Island of Cozumel (*Gaumer*, 1886).

I. Jalapa, (Linn.) Pursh.
In the more open woods, Mascab Pixoy, Jan. 22, 1895 (**148**).

I. Bona-nox, Linn. *Oracion.* **XUTU.**
Climbing over the highest shrubs among the ruins, Chichen Itza, Jan. 20, 1895 (**138** and **123**). In our specimens the calyx is one-eighth the length of the tube of the corolla. Island of Cozumel (*Gaumer*, 1885).

I. puncticulata, Benth.
Completely covering an old well house, base of the Cerro Grande, Izamal, Jan. 13, 1895 (**67**).

I. Jamaicensis (Spreng.) Don.
Convolvulus Jamaicensis, Spreng. Island of Cozumel (*Gaumer*, 1885 and 1886).

I. fastigiata, Sweet
Island of Cozumel (*Gaumer*, 1886). Open woods Mascab Pixoy, Jan. 22, 1895 (**147**).

JACQUEMONTIA, Choisy.

J. pentantha (Jacq.) Don. *Sombaritas azul.* **AK-IXIU.**
Convolvulus pentanthos, Jacq. *Jacquemontia violacea* (Vahl) Choisy. Island of Cozumel (*Gaumer*, 1885 and 1886), open woods near the Cenote of Ix-kil, Chichen Itza, Jan. 20, 1895 (**142**) Mascab Pixoy, Jan. 22, 1895 (**145**).

CONVOLVULUS, Linn.

C. Jamaicensis, Jacq.
Higher banks of the coast, above the clear sand beach, Progreso, Jan. 24, 1895 (**208**). The ovate-lanceolate leaved form.

CUSCUTA, Linn.

C. Americana, Linn.
On weeds in an old Milpa southern end of the Island of Cankun, Jan. 3, 1895. Island of Cozumel (*Gaumer*, 1885 and 1886).

HYDROPHYLLACEAE.

HYDROLEA, Linn.

H. glabra, Choisy.
Yucatan, loc. ignot. (*Johnson*).

NAMA, Linn.

N. Jamaicensis, Linn.
Island of Cozumel (*Gaumer*, 1886).

ASPERIFOLIACEAE.

CORDIA, Linn.

C. dodecandra, DC. *Ciricote.* **COPTÉ.**
Common in gardens. Merida, Progreso, Izamal, and San Miguel, Island of Cozumel; where it is cultivated for its edible fruit. Izamal, Jan. 13, 1895 (**74**). Yucatan, loc. ignot. (*Johnson*). Island of Cozumel (*Gaumer*, 1885).

C. Sebestana, Jacq. *Anachuita.*
Cordia speciosa, R & S. Cultivated at Dolores, Island of Mugeres, for its edible fruit and beautiful flowers. In flower Jan. 3, 1895 (**29**). Also collected here by Dr. Gaumer in 1886.

C. gerascanthoides, Kth. **HABEEM.**
Cordia gerascanthus, Sw. (non Jacq.). A beautiful tree 10 to 30 feet high, with a profusion of pinkish-white flowers. Plentiful about Izamal, especially on the road to Tepakaam, Jan. 15, 1895 (**89**) Island of Mugeres (*Gaumer*, 1886).

C. globosa (Linn.) H.B.K., var. ?
Varronia globosa, Linn. Island of Cozumel (*Gaumer*, 1885).

EHRETIA, Linn.

E. tinifolia, Linn.
Yucatan, loc. ignot. (*Grisebach*).

TOURNEFORTIA, Linn.

T. gnaphalodes (Jacq.) R. Br.
Heliotropium gnaphalodes, Jacq. The principal growth at the inner margin of all the sandy beaches of the Island of Mugeres, North Point, Jan. 1, 1895, (**12**); also collected here by Dr. Gaumer in 1886. Same position at Progreso, and the Island of Cankun.

T. volubilis, Linn.
Island of Mugeres (*Gaumer*, 1886).

HELIOTROPIUM, Linn.

H. inundatum, Sw. **NEMAX.**
Yucatan, loc. ignot. (*Johnson*). Island of Cozumel (*Gaumer*, 1885).

H. parviflorum, Linn. *Rabo de Mico.* **XNA-MAX.**
Campo about Merida, May, 1887. Campo at Chichen Itza, Jan. 18, 1895 (**117**). Yucatan, loc. ignot. (*Johnson*). Island of Cozumel (*Gaumer*, 1885).

VERBENACEAE.

LANTANA, Linn.

L. odorata, Linn.
On rocks inland a few rods from the beach sands, shrubby but kept closely cropped by herbivorous animals. North Point, Island of Mugeres, Jan. 1, 1895 (**10**). Islands of Holbox and Mugeres (*Gaumer*, 1886).

L. Camara, Linn.
Island of Cozumel (*Gaumer*, 1885).

L. involucrata, Linn.
Island of Cozumel (*Gaumer*, 1885).

BOUCHEA, Cham.

B. prismatica (Linn) O. K.
Verbena prismatica, Linn. *Bouchea Ehrenbergii*, Cham. Yucatan, loc. ignot. (*Linden*).

VALERIANODES, Boerh.

E. Jamaicense (Linn) Boerh.
Verbena Jamaicensis, Linn. *Stachytarpheta Jamaicensis*, Vahl. Island of Cozumel (*Gaumer*, 1885). Island of Mugeres (*Gaumer*, 1886).

PRIVA, Adans.

P. lappulacea, (Linn) Pers.
Verbena lappulacea Linn. *Priva echinata*, Juss. Island of Cozumel (*Gaumer*, 1885 and 1886).

PETREA, Linn.

P. volubilis, Jacq. ?
Yucatan, loc. ignot. (*Johnson*).

DURANTA, Linn.

D. repens, Linn.
Duranta Plumieri, Jacq. Island of Cozumel (*Gaumer*, 1885).

VOLKAMERIA, Linn.

V. ligustrina (R. Br.) Jacq.
Clerodendron ligustrinum, R. Br. Yucatan, loc. ignot. (*Johnson*).

LIPPIA, Linn.

L. origanioides, H. B. K.
Yucatan, loc. ignot. (*Johnson*).

L. nodiflora (Linn) Michx.
Verbena nodiflora, Linn. Island of Cozumel along the road to Cedral, Jan. 5, 1895; also at Lagoona Colombia, Jan. 7. Collected here by Dr. Gaumer in 1885, and on the Island of Holbox in 1886.

L. stoechadifolia (Linn.) H.B.K.
Verbena stoechadifolia, Linn. Island of Cozumel (*Gaumer*, 1885). "Not common."

L. ——— sp. (Hemsl.)
Island of Mugeres (*Gaumer*, 1886).

CALLICARPA, Linn.

C. acuminata, H.B.K., var.? (Hemsl.)
Island of Cozumel, (*Gaumer* 1885).

AVICENNIA, Linn.

A. officinalis, Linn.
A. tomentosa, Jacq. Island of Cozumel (*Gaumer*, 1885).

A. nitida, Jacq.
Island of Holbox (*Gaumer*, 1886).

LABIATEAE.

HYPTIS, Jacq.

H. capitata, Jacq.
Yucatan, loc. ignot. (*Johnson*).

H. pectinata, Poit.
Island of Cozumel (*Gaumer*, 1886).

CLINOPODIUM, Linn.

C. Brownei, (Sw.) O. K.
Thymus Brownei, Sw. *Micromeria Brownei*, Benth. In moist soil of a pathway, road to Cedral, Island of Cozumel, Jan. 5, 1895 (**39**). Has the odor of Pennyroyal.

Forma calyce angustiore, Hemsl.
Island of Cozumel (*Gaumer*, 1885).

SALVIA, Linn.

S. occidentalis, Sw. **CAK-NEP.**
Our specimens of the form having mucronate calyx lobes (*S. obscura*, Benth.) Streets of Dolores, Island of Mugeres, Jan. 1, 1895 (**18**). Road to Tepakaam near Izamal, Jan. 14, 1895 (**90**).
Being greatly interested in the similitude of the Maya name Kak-Nep to our Catnep, I finally, upon inquiry, traced it to derivation from a gentleman of New Orleans who visited the Island of Mugeres early in this century; he called the attention of the Indians to the plant saying that it was catnep and very useful in decoction for fever.

S. coccinea, Linn. f. **OIC-XIU.**
This brilliant flowered species is nowhere common, but may be met with here and there throughout the Peninsula. Cerro Grande, Izamal, Jan. 13, 1895. Campo about Chichen Itza, Jan. 18, 1895 (**116**).

S. serotina, Linn.
Island of Cozumel (*Gaumer*, 1885). Island of Mugeres (*Gaumer*, 1886).

TEUCRIUM, Linn.

T. inflatum, Sw.
Along the railroad at Tecantun, Jan. 24, 1895 (**190**), erect ½ to 2½ feet high.

SOLANACEAE.

LYCIUM, Linn.

L. ————sp. (Hemsl.)
Island of Holbox (*Gaumer*, 1886). "Near *L. barbinodum*, Miers. Hemsley."

CAPSICUM, Linn.

C. baccatum, Linn.
Island of Cozumel (*Gaumer*, 1885).

C. frutescens, Linn.
Island of Cozumel (*Gaumer*, 1886).

SOLANUM, Linn.

S. torvum, Sw.
Yucatan, loc. ignot. (*Johnson*).

S. ——— sp. (Hemsl.)
Island of Holbox (*Gaumer*, 1886).

S. Havanense, Jacq. *Yerba mora.* **BA-HA-KAN.**
This shrubby form is quite plentiful about the ruins at Chichen Itza, Jan. 18, 1895 (**120**).

S. verbascifolium, Linn. **TOM-PAAP.**
Among the rocks of the beach, South Point, Island of Mugeres, Jan. 2, 1895 (**25**). Island of Cozumel (*Gaumer*, 1885 and 1886). About the ruined Monjas, Chichen Itza, Jan. 22, 1895 (**140**).

S. nigrum, Linn., var. (Hemsl.)
Island of Cozumel (*Gaumer*, 1885). Island of Mugeres (Gaumer, 1886).

S. Amazonium, Ker.
Island of Cozumel (*Gaumer*, 1885). Along the Tepakaam road near Izamal, Jan. 15, 1895 (**98**).

S. Humboldtii, Willd. **ZUNUMBAK.**
Lycopersicum Humboldtii, Dun. Plentiful about the ruins Chichen Itza, Jan. 20, 1895 (**141**).

DATURA, Linn.

D. Stramonium, Linn. *Chamico.* **XTOHCU.**
Found only near dwellings. Island of Cozumel at Cedral. Waste grounds at Merida, and Mascab Pixoy, Jan. 22, 1895 (**146**). Island of Cozumel (*Gaumer*, 1885).

CESTRUM, Linn.

C. diurnum, Linn.?
Island of Cozumel (*Gaumer*, 1885).

SCROPHULARIACEAE.

RUSSELIA, Jacq.

R. sarmentosa, Jacq.
Island of Cozumel (*Gaumer*, 1885).

MONNIERA, P. Br.

M. Monniera (Linn.) Britton.
Gratiola Monniera, Linn. *Herpestis Monniera*, H.B.K. Island of Cozumel (*Gaumer*, 1885).

M. procumbens (Mill.)
Erinus procumbens, Mill. *Herpestis chamaedryoides*, H.B.K. Island of Cozumel (*Gaumer*, 1885).

CAPRARIA, Linn.

C. biflora, Linn.
Island of Cozumel (*Gaumer*, 1885). Islands of Mugeres and Holbox (*Gaumer*, 1886).

C. biflora pilosa, Griseb. *Clondioso.*
Campo about the ruins, Chichen Itza, Jan 18, 1895, (**118** and **119**).
My **118** agrees with the specimens collected on Fortune and Watling's Islands by Mr. A. S. Hitchcock; and my **119** with those collected by him on the Island of Inagua.

BUECHNERA, Linn.

B. Mexicana, Hemsl. **KABA-CHÉ-CHÉ-BÉ.**
Plentiful in the grass bordering the path at the base of the Great Pyramid, Chichen Itza, Jan. 17, 1895 (**104**).

BIGNONIACEAE.

BIGNONIA, Linn.

B. ——— sp. (Hemsl.)
Island of Cozumel (*Gaumer*, 1885). "Vine 10 to 30 feet; not abundant. No fruit, and only imperfect leaves." Hemsley.

B ——— sp. (Hemsl.)
Island of Cozumel (*Gaumer*, 1885). "Vine 10 to 100 feet; no fruit, and but imperfect leaves." Hemsley.

TECOMA, Linn.

T. stans (Linn.) Juss. *Flor Amarillo.* **XKANLOL.**
Bignonia stans, Linn. On the summit of the Cerro Grande Izamal, Jan. 13, 1895 (**55**), in flower and fruit. At the foot of the Great Pyramid, Chichen Itza, Jan. 18, 1895 (**124**), 6 to 8 feet high.

CRESCENTIA, Linn.

C. Cujete, Linn. *Cujete.* **LUCH.**
One specimen only of this tree observed, that at Mascab Pixoy. It was out of season both as to flower and fruit.

PARMENTIERA, DC.

P. edulis (Mocq.) DC. *Pepina.*
Cultivated ? Street in Dolores, Island of Mugeres, Jan. 2, 1895 (**30**). A small umbrageous tree 15 to 25 feet high, with a strict cylindrical trunk. Fruits 6 to 10 inches long.

PEDALIACEAE.

MARTYNIA, Linn.

M. diandra, Glox.
Martynia angulosa, Lam. Yucatan, loc. ignot. (*Johnson*).

ACANTHACEAE.

TUBIFLORA, Gm.

T. squamosa (Jacq.) O.K.
Verbena squamosa, Jacq., *Elytraria tridentata*, Vahl. Campo about Izamal, Jan. 23, 1895 (**188**).

RUELLIA, Linn.

R. paniculata, Linn. **KABAUCHÉ, ɔAKALBAK.**
A low straggling shrub. On the slopes of the Cerro Grande Izamal, Jan. 13, 1885 (**57**). Margin of the scrub along the Tepakaam road near Izamal, Jan. 14, 1895 (**93**). Campo about the ruins Chichen Itza, Jan. 17, 1895 (**105** and **139**).
The Maya ɔakalbak is applied to all Ruellias of this form, the word Kabauché meaning really nothing more distinctive than "a small bush."

BLECHUM, P. Br.

B. Brownei, Juss.
Campo about Izamal, Jan. 23, 1895 (**171**).

B. Brownei *var.* (Hemsl.)
Island of Cozumel (*Gaumer* 1885).

BRAVAISIA, DC.

B. tubiflora, Hemsl.
Type locality, Island of Cozumel (*Gaumer* 52, 1885); "Tree 25 feet; not abundant." Higher banks of sea shore Progreso, Jan. 26, 1895 (**206**). Low shrub, flowers white with a tinge of purple, lower lip purple, throat yellow, buds yellowish white.

APHELANDRA, R. Br.

A. pectinata, Nees.
Shrub 15 feet high. Tepakaam road near Izamal, Jan. 14, 1895 (**91**).

DIAPEDIUM, Koenig.

D. assurgens (Linn). O.K.
Justicia assurgens, Linn. *Dicliptra assurgens* Juss. Yucatan loc. ignot. (*Johnson.*) Island of Cozumel (*Gaumer*, 1885 and 1886).

DIANTHERA, Gronov.

D. comata, Linn.
Yucatan loc. ignot. (*Johnson*).

TETRAMERIUM, Nees.

T. hispidum, Nees.
Edges of Copses, Chal-tun-ha road near Izamal, Jan. 14, 1895 (**78**).

T. costatum (Gray).
Henrya costata Gray. Among the rocks on the Cerro Grande Izamal, Jan. 13, 1895 (**65**).

DIANTHERA, Gronov.

D. sessilis, Griseb.
In the scrub along the Chal-tun-ha road near Izamal, Jan. 14, 1895 (**79**).

PLANTAGINACEAE.

PLANTAGO, Linn.

P. major, Linn.
Island of Cozumel (*Gaumer*, 1885).

RUBIACEAE.

RACHICALLIS, DC.

R. Americana (Jacq.) O. K.
Hedysotis Americana, Jacq. *Rachicallis rupestris* (Sw.) DC. Island of Cozumel (*Gaumer*, 1885).

RONDELETIA, Linn.

R. stenosiphon, Hemsl.
Yucatan, loc. ignot. (*Johnson*).

COUTAREA, Aubl.

C. octomera, Hemsl.
Type locality, Island of Cozumel (*Gaumer*, 1885). "Very rare shrub, flowers greenish-yellow."

RANDIA, Linn.

R. aculeata, Linn.
Island of Cozumel (*Gaumer*, 1885).

R. latifolia, Linn.
Island of Cozumel (*Gaumer*, 1885).

R. ? longiloba, Hemsl.
Type locality, Island of Cozumel (*Gaumer*, 1885). "Tree 25 feet; very common and very aromatic."

HAMELIA, Jacq.

H. patens, Jacq. **XKAN-NAN.**

Shrub 6 to 8 feet high, growing upon the lower terrace of the ruined Monjas, Chichen Itza, Jan. 1895 (**122**). Island of Cozumel (*Gaumer*, 1885 and 1886).

H. chrysantha, Sw.

Along the roadside near Cedral, Island or Cozumel, Jan. 5, 1895 (**41**).

GUETTARDA, Linn.

G. scabra (Linn.) Lam.

Mathiola scabra, Linn. Yucatan, loc. ignot. (*Linden*).

MACHAONIA, Humb. & Bonpl.

M. Lindeniana, Baill.

Yucatan, type loc. ignot. (*Linden*).

ERITHALIS, P. Br.

E. fruticosa, Linn.

A low shrub, 4 to 5 feet high, forming a part of the general littoral growth at the margin of the sandy beach, North Point, Island of Mugeres, Jan. 2, 1895 (**27**). Islands of Holbox and Mugeres (*Gaumer*, 1886).

CHIOCOCCA, P. Br.

C. racemosa, Jacq.

Island of Cozumel (*Gaumer*, 1885), "varietates." Hemsl.

ASEMNANTHA, Hook f.

A. pubescens, Hook f.

Humid forests, loc. ignot. (*Linden*).

STRUMPFIA, Jacq.

S. maritima, Jacq.

Island of Mugeres, on the higher reef rocks almost in constant spray, Jan. 1, 1895 (**11**). Collected here also by Dr. Gaumer in 1886.

A low, spreading, shrubby growth resembling a Juniper or a frutescent Erica. This species seems to have reached this Island from the Bahamas without, as far as I can learn, finding a foothold upon any of the intervening islands.

COUSSAREA, Aubl.

C. speciosa, Aubl. ?

Yucatan, loc. ignot. (*Linden*).

MORINDA, Linn.

M. Royoc, Linn.
Island of Cozumel (*Gaumer*, 1885). "A littoral bush 1 to 3 feet; abundant."

ERNODEA, Sw.

E. littoralis, Sw.
On the higher beach above the clear beach sands, Progreso, Jan. 26, 1895 (**207**).
Shrubby. Flower tube deep brownish-lake externally, white within corolla, teeth sharply recurved.
Island of Cozumel (*Gaumer*, 1885). "Bush 1 to 3 feet; abundant" Islands of Holbox and Mugeres (*Gaumer*, 1886).

BORRERIA, G. F. W. Mey.

B. thymifolia, Griseb.
On the Cerro Grande, Izamal, Jan. 13, 1895 (**63**).

SPERMACOCE, Linn.

S. laevis, Lam.
Yucatan, loc. ignot. (*Johnson*).

S. verticillata, Sw.
Island of Holbox (*Gaumer*, 1886).

S. teunior, Lam (not Linn.)
Campo at Izamal, Jan 23, 1895 (**183**). Island of Cozumel (*Gaumer*, 1885 and 1886).

S. Haenkeana, Hemsl. *Flor de Manzanillo do Campo.*
Campo about the ruins, Chichen Itza, Jan. 17, 1895 (**113**). My specimens agree with Pringle's 2453 from Jalisco, except that the glomerules of mine are somewhat smaller.
The Mayas claim that this form has severe ptarmacoid properties.

TRIODON, DC

T. angulatum, Benth.
Yucatan, loc. ignot. (*Johnson*).

CAPRIFOLIACEAE.

SAMBUCUS, Linn.

S. Mexicana, Presl.
About the ruins at the southern end of the Island of Cankun, Jan. 3, 1895, not in bloom. Island of Cozumel (*Gaumer*, 1885).

CUCURBITACEAE.

MELOTHRIA, Linn.

M. pendula, Linn. **XTULU.**

About the ruined Monjas, Chichen Itza, Jan. 17, 1895 (**109** and **110**). Very young but in flower.

The Maya name Xtulu "Little Watermelon" is applied to various cucurbits.

SICYDIUM, Schlecht.

S. tamnifolium, (Poir.) Cogn.

At Tiop, (*Linden*). Yucatan, loc. ignot. (*Schott.*)

S. ——— sp.? (Hemsl.)

Island of Cozumel (*Gaumer*, 1885). "Vine 15 feet; rare. Foliage and male flowers only." Hemsley.

MOMORDICA, Linn.

M. Charantia, Linn.

Island of Cozumel (*Gaumer*, 1885).

LUFFA, Linn.

L. cylindrica (Linn.) Roem.

Momordica cylindrica, Linn. Yucatan, loc. ignot. (*Schott*).

CAYAPONIA, Manso.

C. alata, Cogn.

Type locality, the Hacienda Sara-Grossa (Schott).

C. grandiflora, Cogn.

Type locality, Picul (*Schott*).

CAMPANULACEAE.

ISOTOMA, Lindl.

I. longiflora (Linn.) Presl.

Lobelia longiflora, Linn. Island of Cozumel (*Gaumer*, 1885). "Not common."

GOODENIACEAE.

SCAEVOLA, Linn.

S. Plumieri (Linn.) Vahl.

Lobelia Plumieri, Linn. Islands of Holbox and Mugeres (*Gaumer*, 1886).

COMPOSITAE.*

VERNONIA, Schreb.

V. lanceolaris, DC.

Yucatan, loc. ignot. (*Johnson*).

V. Schiedeana, Less.

Yucatan, loc. ignot. (*Johnson*).

* My numbers were kindly determined by Professor J. M. Coulter.

OLIGANTHES, Cass.

O. oxylepis, Benth & Hook.
Yucatan, loc. ignot. (*Johnson*).

ALOMIA, H.B.K.

A. ageratoides, H.B.K. ?
Ethulia ageratoides, Spreng. Island of Cozumel (*Gaumer*, 1885). "Differs from the next species only in having white flowers and epappose achenes." Hemsley.

AGERATUM, Linn.

A. intermedium, Hemsl.
Type locality, Island of Cozumel (*Gaumer*, 1885). "Abundant."

A. corymbosum, Zucc.
Campo about Izamal, Jan. 13, 1895, (**72**).

EUPATORIUM, Linn.

E. odoratum, Linn.
Yucatan, loc. ignot. (*Johnson*).

E. ——— sp. (Hemsley).
Island of Cozumel (*Gaumer*, 1885). "Rare, flowers white tinged with dull yellow."
Professor Hemsley says: "This agrees with a specimen from Yucatan and Tabasco (Torrey 1850) together with which it may be or may not be an undescribed species." The Yucatan and Tabasco species is doubtless one of the Johnson collection, as Professor Torrey contributed that collection to Kew.

E. populifolium, H.B.K.
Yucatan, loc. ignot. (*Johnson*).

E. paniculatum, Schrad.
Island of Cozumel (*Gaumer*, 1886).

E. ——— sp. (Hemsl.)
Island of Cozumel (*Gaumer*, 1886), "Near E. conyzoides, Vahl."

E. conyzoides, Vahl.
Campo about the ruins, Chichen Itza, Jan. 17, 1895 (**115**).

WILLOUGHBIA, Neck.

W. Houstonis (Linn.) O.K.
Eupatorium Houstonis, Linn. *Mikania Houstonis*, Willd. Yucatan, loc. ignot. (*Johnson*).

CHRYSOPSIS, Nutt.

C. ——— sp.
Roadside near Izamal, Jan. 15, 1895 (**97**). Three feet high, nearly out of flower.

EGLETES, Less.

E. viscosa, Less.
Yucatan, loc. ignot. (*Johnson*).

AMELLUS, P. Br.

A. aspera (Jacq.) O.K.
Calea aspera, Jacq. *Melanthera deltoidea*, Michx. Island of Cozumel (*Gaumer*, 1886).

A. niveus (Linn.) O.K.
Bidens nivea, Linn. *Melanthera hastata*, Rich. On the beach near the Lagoona Colombia, Island of Cozumel, Jan. 7, 1895 (**51**). Island of Cozumel (*Gaumer*, 1885). Yucatan, loc. ignot. (*Johnson*).

ERIGERON, Linn.

E. Canadensis, Linn.
Island of Cozumel (*Gaumer*, 1885).

PLUCHEA, Cass.

P. purpurascens (Sw.) DC.
Conyza purpurascens, Sw. Island of Cozumel (*Gaumer*, 1885).

P. odorata (Linn.) Cass.
Conyza odorata, Linn. Streets of Izamal, Jan. 23, 1895 (**176**). Island of Mugeres (*Gaumer*, 1886).

PARTHENIUM, Linn.

P. Hysterophorus, Linn.
Island of Cozumel (*Gaumer*, 1885 and 1886).

AMBROSIA, Linn.

A. hispida, Pursh.
Ambrosia crithmifolia, DC. Sea shore, Island of Cozumel (*Gaumer*, 1885). "Very Abundant."

BLAINVILLEA, Cass.

B. dichotoma (Murr.) Cass.
Verbesina dichotoma, Murr. Island of Cozumel (*Gaumer*, 1886).

BORRICHIA, Adans.

B. arborescens (Linn). DC.
Buphthalmium arborescens, Linn. Sea shore, Island of Cozumel (*Gaumer*, 1885). "Bushy, 3 feet; abundant." Islands of Mugeres and Holbox (*Gaumer*, 1886).

WEDELIA, Jacq.

W. trilobata (Linn.) Hitch.
Silphium trilobatum, Linn. *Wedelia carnosa*, Pers. Island of Cozumel (*Gaumer*, 1885).

ASPILIA, Thou.

A. ——— sp.
Old clearing near Mascab Pixoy, Jan. 22, 1895 (**154**).

VIGUIERA, Less.

V. helianthoides, H.B.K.
Island of Cozumel (*Gaumer*, 1886).

ELEUTHERANTHERA, Poit.

E. divaricata (Rich.)
Dysodium divaricatum, Rich. *Melampodium divaricatum*, DC. Patio residence of Dr. Gaumer, Izamal, Jan. 13, 1895 (**71**). Island of Cozumel (*Gaumer*, 1885 and 1886).

SPILANTHES, Jacq.

S. repens (Walt.) Michx.
Anthemis repens, Walt. Island of Cozumel, marshy places (*Gaumer*, 1885 and 1886). "Not abundant."

S. beccabunga, DC.
Moist soil of pathway to Cedral, Island of Cozumel, Jan. 5, 1895 (**43**).

HELIANTHELLA, T. & G.

H. ——— sp. **TAH.**
Plentiful in the clearing at the base of the Great Pyramid, Chichen Itza, Jan. 17, 1895 (**103**), 4 to 6 feet high, flowers bright King's yellow.

ZEXMENIA, Llav. & Lex.

Z. trachylepes, Hemsl.
Yucatan, type loc. ignot. (*Johnson*).

SYNEDRELLA, Gaertn.

S. nodiflora, Gaertn.
Campo about Izamal, Jan. 23, 1895 (**185**).

BIDENS, Linn.

B. pilosa, Linn.
Island of Cozumel (*Gaumer*, 1885).

B. Alansensis, H.B.K.
Compo about Izamal, Jan. 23, 1895 (**170**).

CALEA, Linn.

C. ——— sp.
Old clearing near Mascab Pixoy, Jan. 22, 1895 (**153**).

TRIDAX, Linn.

T. procumbens, Linn.
Campo about Izamal, Jan. 23, 1895 (**218**). Yucatan, loc. ignot. (*Linden*). Island of Cozumel (*Gaumer*, 1885).

FLAVERIA, Juss.

F. linearis, Lag.
Island of Cozumel (*Gaumer*, 1885).

F. longifolia, A. Gray.
Island of Holbox (*Gaumer*, 1886).

TAGETES, Linn.

T. tenuifolia, Cav.
Campo about Izamal, Jan. 23, 1895 (**179**).

NEUROLAENA, R. Br.

N. lobata (Linn.) R. Br.
Conyza lobata, Linn. Yucatan, loc. ignot. (*Johnson*).

TRIXIS, P. Br.

T. radialis (Linn.) O.K.
Perdicium radialis, Linn. *Trixis Inula*, Crantz, *Trixis frutescens* R. Br. Apex of the Cerro Grande, Izamal, Jan. 13, 1895 (**56**).

ORDERS.	Genera.	Species.	Mainland.	Insular Species.				Dr. Gaumer.	Author.	Johnson.	Linden.	Schott.
				Cozumel.	Mugeres.	Holbox.	Cankun.					
Acanthaceæ	9	11	10	3	..	..	..	3	8	2	..	..
Amaranthaceæ	6	8	5	2	1	1	..	4	4	1	..	..
Amaryllidaceæ	1	1	1	1	..	..	..	1	..	1	..	..
Anacardiaceæ	3	4	1	4	..	..	..	3	1	1	..	..
Anonaceæ	1	3	2	1	..	..	..	1	2	..	..	..
Apocynaceæ	5	8	4	2	1	..	..	4	2	2	..	..
Araceæ	1	1	1	..	..	..	..	..	1	..	..	..
Aristolochiaceæ	1	1	1	..	..	..	..	..	1	..	..	..
Asclepiadaceæ	6	7	5	2	1	1	..	3	2	..	3	..
Asperifoliaceæ	4	9	6	4	4	..	1	7	5	2	..	..
Bignoniaceæ	4	5	2	2	1	..	..	2	3	..	..	..
Bixaceæ	3	3	3	..	..	..	..	..	2	1	..	..
Bombacaceæ	3	5	5	..	..	..	..	..	4	1	..	..
Bromeliaceæ	5	8	6	1	..	1	..	1	2	3	..	3
Burseraceæ	1	1	1	1	1	..	..	..	1	..	..	..
Cactaceæ	2	4	1	2	3	..	..	1	3	..	..	..
Campanulaceæ	1	1	..	1	..	..	..	1	..	..	..	..
Cannaceæ	1	1	..	1	..	..	..	1	..	..	..	..
Capparidaceæ	2	2	2	..	..	..	..	..	1	..	1	..
Caprifoliaceæ	1	1	..	1	..	..	1	1	1	..	..	..
Caricaceæ	2	2	2	1	1	..	1	1	2	..	..	..
Casuarinaceæ	1	1	1	1	1	..	1	..	1	..	..	..
Celastraceæ	2	2	..	1	1	..	..	2	..	..	..	..
Chenopodiaceæ	2	2	..	..	..	2	..	2	..	..	..	..
Combretaceæ	2	2	..	2	..	1	..	2	..	..	..	..
Commelynaceæ	2	3	1	2	1	..	..	2	..	..	..	1
Compositæ	30	42	24	21	2	2	..	22	15	11	..	..
Convolvulaceæ	5	13	7	9	..	..	1	9	7	1	..	..
Crassulaceæ	1	1	1	..	..	..	..	..	1	..	..	..
Cruciferæ	1	1	1	1	1	..	..	1	1	..	..	..
Cucurbitaceæ	5	7	5	1	..	..	..	2	1	..	1	4
Cycadaceæ	1	1	1	..	..	..	..	..	..	..	..	..
Cyperaceæ	2	6	1	3	2	1	..	4	1	..	..	1
Dioscoreaceæ	1	2	1	1	..	..	..	1	..	..	1	..
Euphorbiaceæ	6	22	13	13	5	2	1	11	15	..	1	..
Filices	8	9	4	5	..	..	..	3	6	..	..	..
Gentianaceæ	1	1	..	1	..	1	..	1	..	..	..	..
Goodeniaceæ	1	1	..	..	1	1	..	1	..	..	..	..
Gramineæ	8	11	5	4	1	4	..	5	5	..	1	..
Guttiferæ	1	1	..	1	..	..	..	1	..	..	..	..
Hepaticæ	3	4	2	2	..	..	..	..	4	..	..	..
Hippocrateaceæ	1	1	..	1	..	.	..	1	..	..	..	..
Hydrophyllaceæ	2	2	1	1	..	..	..	1	..	1	..	..
Iridaceæ	1	1	1	..	..	..	..	..	..	..	1	..
Labiateæ	4	7	4	3	2	.	..	3	4	1	..	..
Lauraceæ	3	3	1	1	..	1	..	2	1	..	..	..
Leguminosæ	32	58	31	30	9	..	..	33	18	13	4	..
Liliaceæ	2	2	2	..	..	..	..	..	..	..	..	2
Loasaceæ	1	1	1	..	..	..	..	..	1	..	..	..
Lythraceæ	1	1	1	1	..	..	..	1	..	1	..	..
Malpighiaceæ	5	8	5	3	..	..	..	3	..	4	1	..
Malvaceæ	6	10	8	2	..	..	..	2	7	1	..	..
Melastomaceæ	3	3	3	..	..	..	..	..	..	3	..	..
Meliaceæ	3	4	2	2	..	..	..	2	1	..	1	..
Moraceæ	2	4	4	1	1	..	..	..	4	..	..	..
Musaceæ	1	2	2	2	1	..	2	..	2	..	..	..

ORDERS.	Genera.	Species.	Mainland.	Insular Species.				Dr. Gaumer.	Author.	Johnson.	Linden.	Schott.
				Cozumel.	Mugeres.	Holbox.	Cankun.					
Musci	3	3	2	2	..	..	..	..	3	..	..	..
Myrsinaceæ	2	3	1	2	..	1	..	2	..	1	..	..
Myrtaceæ	3	3	2	1	..	..	..	2	2	..	..	..
Nyctaginaceæ	2	3	3	1	..	..	..	1	3	..	..	..
Nymphaceæ	2	2	2	..	..	..	..	..	1	..	1	..
Orchidaceæ	1	1	1	..	..	..	..	..	1	..	..	..
Palmeæ	3	4	2	3	4	..	3	2	3	..	..	..
Pandaneæ	1	2	2	..	..	..	..	..	2	..	..	..
Papaveraceæ	1	1	1	1	..	..	..	1	1	..	..	..
Passifloraceæ	1	2	..	2	1	..	..	2	..	..	..	..
Pedaliaceæ	1	1	1	..	..	..	..	..	..	1	..	..
Phytolaccaceæ	2	5	3	2	2	..	..	2	1	2	1	..
Piperaceæ	1	1	1	..	..	..	..	..	..	..	1	..
Plantaginaceæ	1	1	..	1	..	..	..	1	..	..	..	..
Plumbaginaceæ	1	1	1	..	..	..	..	..	1	..	..	..
Polygalaceæ	1	1	1	..	..	..	..	..	..	1	..	..
Polygonaceæ	2	3	..	3	1	..	..	3	1	..	..	..
Ranunculaceæ	1	3	3	1	..	..	..	1	1	1	..	1
Rhamnaceæ	1	1	1	..	..	..	..	..	..	..	1	..
Rhizophoraceæ	1	1	..	1	..	1	1	1	1	..	..	..
Rubiaceæ	17	23	12	11	3	3	..	14	8	3	4	..
Rutaceæ	6	9	4	4	2	..	..	4	5	..	..	..
Sapindaceæ	5	7	5	2	..	..	..	2	2	3	1	..
Sapotaceæ	3	3	2	1	1	..	..	1	2	..	..	..
Scrophulariaceæ	4	6	2	4	1	1	..	4	2	..	..	..
Simarubaceæ	2	2	1	1	1	1	..	2	1	..	..	..
Smilaceæ	1	1	..	1	..	..	..	1	..	..	..	..
Solanaceæ	5	12	5	7	2	2	..	9	5	1	..	..
Sterculiaceæ	6	7	5	4	..	..	..	3	3	2	1	..
Tiliaceæ	1	2	2	..	..	..	..	..	..	2	..	..
Urticaceæ	1	1	1	..	..	..	..	..	..	1	..	..
Verbenaceæ	10	16	4	9	3	3	..	12	2	3	1	..
Violaceæ	1	1	1	..	..	..	..	..	..	1	..	..
Vitaceæ	1	2	1	1	..	..	..	1	..	1	..	..
Zygophyllaceæ	1	2	1	1	..	..	..	1	1	..	..	..
Totals: Orders, 91...	307	471	264	210	62	30	12	224	192	75	25	12

SUMMARY.

Mainland Flora..... 264 species.

Insular Flora { Cozumel ..... 210 / Sand Islands..... 104 } 314 "

Index.

PAGE.

Abutilon 29
Acacia 19
Acalypha 27
ACANTHACÆ 46
Achyranthes 15
Achras 37
Adiantum 8
Æchmea 12
Æschynomene 22
Ageratum 51
Alicastrum 14
Alomia 51
Alternanthera 16
Alvaradoa 33
AMARANTHACEÆ 15
Amaranthus 15
AMARYLLIDACEÆ 12
Amaryllis 12
Ambrosia 52
Amellus 52
Amerimnon 22
ANACARDIACEÆ 32
Ananas 12
Aneimia 9
Anoda 29
Anona 17
ANONACEÆ 17
Anthemis 53
Antheropogon 10
Antigonon 15
Aphelandra 46
APOCYNACEÆ 37
ARACEAE 11
Argemone 18
Aristolochia 15
ARISTOLOCHIACEÆ 15
Arum 11
ASCLEPIADACEÆ 38
Asclepias 38
Asemnantha 48
ASPERIFOLIACEÆ 40
Aspidium 9

PAGE.

Aspilia 53
Asplenium 8
Ateleia 21
Atriplex 15
Avicennia 42
Ayenia 31
Barbula 8
Bauhinia 20
Bidens 54
Bidens 52
Bignonia 45
Bignonia 45
BIGNONIACEÆ 45
Bixia 32
BIXACEÆ 31
Blainvillea 52
Blechum 46
Bœrhaavia 17
BOMBACACEÆ 30
Bombax 30
Bombax 30
Borreria 49
Borrichia 53
Bouchea 41
Bouteloua 10
Bradburya 23
Bravaisia 46
Bromelia 11
BROMELIACEAE 11
Brosimum 14
Bryophyllum 18
Buceras 36
Buechnera 45
Bumelia 37
Bunchosia 24
Buphthalmium 53
Bursera 25
BURSERACEÆ 25
Byrsonima 24
Cabomba 17
CACTACEÆ 35
Cactus 35

	PAGE.
Cæsalpinia	21
Cakile	18
Calea	54
Calea	52
Calliandra	19
Callicarpa	42
Callisia	13
Calopogonium	23
Calyptrion	32
CAMPANULACEÆ	50
Canna	13
CANNACEÆ	13
CAPPARIDACEÆ	18
Capparis	18
Capraria	45
CAPRIFOLIACEÆ	49
Capsicum	43
Carica	34
CARICACEÆ	34
Carolinea	30
Cassia	20
Cassine	32
Cassytha	18
Castalia	17
Casuarina	14
CASUARINACEÆ	14
Cattleya	13
Cayaponia	50
Cedrela	26
Ceiba	30
CELASTRACEÆ	32
Celosia	15
Cenchrus	9
Centrosema	23
Cereus	35
Cestrum	44
Chalcas	25
Cheilanthes	8
Cheilolejunea	7
CHENOPODIACEÆ	15
Chiococca	48
Chloris	10
Chrysobalanus	19
Chrysopsis	52
Citrus	25
Clematis	17
Clerodendron	42
Clidemia	36
Clinopodium	43
Clitoria	23

	PAGE.
Coccoloba	15
Cochlospermum	31
Cocos	11
Coix	9
Conocarpus	36
Conostegia	36
CONVOLVULACEÆ	39
Convolvulus	40
Convolvulus	39
Conyza	52, 54
COMBRETACEÆ	36
Commelina	13
COMMELYNACEÆ	13
COMPOSITÆ	50
Corchoras	34
Cordia	40
Corynostylis	32
Cotyledon	18
Coussarea	48
Coutarea	47
Cracca	22
CRASSULACEÆ	18
Cratæva	18
Crescentia	45
Crossopetalum	32
Crotalaria	21
Croton	26
CRUCIFERÆ	18
CUCURBITACEÆ	50
Cuscuta	40
CYCADACEÆ	9
Cynanchum	38
CYPERACEÆ	10
Cyperus	10
Dalbergia	22
Dalea	21
Dasylirion	12
Datura	44
Desmanthus	20
Desmodium	22
Dianthera	47
Dianthera	47
Diapedium	46
Dicliptra	46
Dioon	9
Dioscorea	13
DIOSCOREACEÆ	13
Diphysa	22
Dolichos	23
Dolichos	23

	PAGE.
Duranta	42
Dysodium	53
Egletes	52
Ehretia	40
Elæodendron	32
Eleusine	10
Eleutheranthera	53
Elytraria	46
Ensenbeckia	24
Enslenia	38
Eragrostis	10
Erigeron	52
Erinus	44
Eriodendron	30
Erithalis	48
Ernodea	49
Erythrina	23
Erythrina	22
Ethulia	51
Eugeina	36
Eulejunea	7
Eupatorium	51
Eupatorium	51
Euphorbia	27
EUPHORBIACEÆ	26
Eustoma	37
Evolvulus	39
Ficus	14
Fimbristylis	11
Flaveria	54
Glactia	23
Galega	22
GENTIANACEÆ	37
Gentiania	37
Gliricidia	22
Gomphræna	16
GOODENIACEÆ	50
Gouania	35
GRAMINEÆ	9
Gratiola	44
Gronovia	35
Guazuma	31
Guettarda	48
Guilandina	21
GUTTIFERÆ	31
Hæmatoxylon	21
Hamelia	48
Hechtia	12
Hedysarum	22
Hedysotis	47

	PAGE.
Helianthella	53
Heliotropium	41
Heliotropium	41
Hemitelia	8
Henrya	47
HEPATICÆ	7
Herpestris	44
Heteropterys	24
Hibiscus	30
Hippeastrum	12
Hippocratea	33
HIPPOCRATEACEÆ	33
Hiræa	23
Hisingera	32
Hydrolea	40
HYDROPHYLLACEÆ	40
Hyptis	42
Ichthyomethia	22
Indigofera	21
Ipomœa	39
Iria	11
IRIDACEÆ	13
Irisine	16
Isotoma	50
Jacaratia	35
Jacquemontia	39
Jacquinia	36
Jatropha	27
Justicia	46
LABIATEÆ	42
Lantana	41
LAURACEÆ	17
Lawsonia	35
LEGUMINOSÆ	19
Lerchea	15
Leucæna	19
LILIACEÆ	12
Lippia	42
LOASACEÆ	35
Lobelia	50
Lonchocarpus	22
Lucuma	37
Luffa	50
Lycium	43
Lycopersicum	44
Lysiloma	19
LYTHRACEÆ	35
Machaonia	48
Macroscepis	38
Malpighia	24

	PAGE.
MALPIGHIACEÆ	23
Malva	29
MALVACEÆ	29
Malvastrum	29
Malvaviscus	30
Malveopsis	29
Mammea	31
Mangifera	32
Marsdenia	39
Martynia	46
Mastigolejunea	7
Matiola	48
Maxmilianea	31
Megiphanes	16
Meibomia	22
Melampodium	53
Melanthera	52
Melastoma	36
MELASTOMACEÆ	36
Melia	26
MELIACEÆ	26
Melochia	31
Metastelma	38
Melothria	50
Micromeria	43
Mikania	51
Mimosa	20
Mimosa	19
Mirabilis	16
Momordica	50
Momordica	50
Monniera	44
MORACEÆ	14
Morinda	49
Murraya	25
Musa	13
MUSACEÆ	13
MUSCI	7
Myginda	32
Myroxylon	32
MYRSINACEÆ	36
MYRTACEÆ	36
Myrtus	36
Nama	40
Nectandra	18
Neurolæna	54
NYCTAGINACEÆ	16
Nymphæa	17
NYMPHACEÆ	17
Oliganthes	51
Opuntia	35
ORCHIDACEÆ	13
Pachira	30
PALMÆ	11
PANDANACEÆ	9
Pandanus	9
Panicum	9
PAPAVERACEÆ	18
Parathesis	37
Parmentiera	45
Parosela	21
Parthenium	52
Passiflora	34
PASSIFLORACEÆ	34
Paulinia	33
Paulinia	33
PEDALIACEÆ	46
Perdicium	54
Persea	17
Petiveria	16
Petrea	42
Phaseolus	23
Philibertia	38
Philodendron	11
Phyllanthus	26
PHYTOLACCACEÆ	16
Picramnia	25
Pimenta	36
Piper	13
PIPERACEÆ	13
Piscidia	22
Pithecolobium	19
PLANTAGINACEÆ	47
Plantago	47
Pluchea	52
Pleroma	36
PLUMBAGINEÆ	37
Plumbago	37
Plumeria	38
Poinciana	21
Polygala	26
POLYGALACEÆ	26
POLYGONACEÆ	15
Polypodium	7
Priva	41
Psidium	36
Ptelea	25
PTERIDOPHYTA	8
Pteris	8
Pterocarpus	21

	PAGE.
Rachicallis	47
Randia	47
RANUNCULACEÆ	17
Rauwolfia	37
RHAMNACEÆ	33
Rhexia	36
Rhizophora	35
RHIZOPHORACEÆ	35
Rhus	32
Rhynchosia	23
Ricinus	27
Rivina	16
Rondeletia	47
ROSACEÆ	19
Roulinia	38
RUBIACEÆ	47
Ruellia	46
Russelia	44
Rutacearum	24
RUTACEÆ	24
Sabal	11
Salvia	43
Sambucus	49
SAPINDACEÆ	33
Sapindus	33
SAPOTACEÆ	37
Sarcostemma	38
Scævola	50
Schmidelia	33
SCROPHULARIRCEÆ	44
Seriana	33
Serjania	33
Sicydium	50
Sida	29
Sida	29
Silphium	53
SIMARUBACEÆ	25
SMILACEÆ	12
Smilax	12
SOLANACEÆ	43
Solanum	44
Sophora	21
Spermacoce	49
Spilanthes	53
Spondias	32
Stachytarpheta	41
Stenotaphrum	10
Sterculia	31
STERCULIACEÆ	31
Stereophyllum	7
Strumpfia	48
Suriana	25
Synedrella	54
Tabernæmontana	38
Tagetes	54
Tamarindus	20
Tecoma	45
Tephrosia	22
Terminalia	36
Tetramerium	47
Teucrium	43
Theobroma	31
Thevetia	38
Thrinax	11
Thymus	43
Tibouchina	36
Tigridia	13
TILIACEÆ	34
Tillandsia	12
Tournefortia	41
Tribulus	24
Trichilia	26
Tridax	54
Triodon	49
Triumfetta	34
Trixis	54
Tubiflora	46
Urera	14
URTICACEÆ	14
Valerinodes	41
Vallesia	37
Verbena	41, 42, 46
VERBENACEÆ	41
Verbesina	52
Vernonia	50
Viguiera	53
Villamilla	16
Viola	32
VIOLACEÆ	32
VITACEÆ	33
Vitis	33
Volkameria	42
Waltheria	31
Wedelia	53
Willoughbia	51
Xylosma	32
Yucca	12
Zanthoxylum	24
Zexmenia	53
ZYGOPHYLLACEÆ	24

Local Names.

	PAGE.
Abal	32
Abrojo	24
Aguacate	17
AK-IXIU	39
AKUM	14
Alamo	14
Amapola blanca	30
Amapola colorado	30
Amapolite	29
Anachuita	40
Arnatto	32
BA-HA-KAN	44
Bayal	11
BEL-CINICHE	33
Bonete	35
BOX-HAAS	13
BUUL	23
Cacao	31
Cajera	25
CAK-NEP	43
CANCHOH	35
Cardosanto	18
Cedro	26
Ceiba	30
CHACAH	25
CHACHAL-HAAS	37
Chamico	44
CHANXNUC	24
CHE-CHE-BE	29
CHICH-PUT	34
Chicle	37
CHICAN	23
CHOM	11
CHUN	31
Ciricote	40
Clondioso	45
Coco	11
Copo	14
COPTE	40
CUCHILXIN	38
Cujete	45
Flor Amarillo	45

	PAGE.
Flor de Manzanillo do Campo	49
Frijol	23
Guayabo	36
HABEEM	40
Higuerilla	27
HOBONKAK	29
HOL	30
KABA-CHE-CHE-BE	45
KABANCHE	46
KUCHE	26
KULCHÉ	26
KUMCHÉ	35
KUYCHÉ	30
LAB	7
LAL-MUCH	35
Laurel de la India	14
Limon	25
Limoncillo	25
LUCH	45
Mamey Colorado	37
Mango	32
Naranja	25
NEMAX	41
Nopal	35
ON	17
Oracion	39
OX	14
PAH	9
PAKAN	35
Papaya	34
Papaya los Pajaros	34
Pata de Vaca	20
PEEM	30
Pepina	45
Picar	27
PICHI	36
Pina	12
Pitaya	35
PIXOY	31
Pochote	30
PUT	34
Rabo de Mico	41

The Maya names in small capitals.

	PAGE.
Ramon	14
Sabacnicté	38
Sac-hass	13
Sac-piché	15
Sac-xieu	29
Sapote	37
Saramuyo	17
Sombaritas azul	39
Sorillo blanco	15
Sutz-pakal	25
Tah	53
Tapché	35
Tauch-ya	37
Tom-paap	44
Tuna	35
Ɔakalcak	46
Ɔalmuy	17
Ɔic-xiu	43
Ɔipil	9
Uba	15
Urucu	32
Uɔupek	38
Xaan	11
Xcuyché	30
Xkanlol	45
Xkan-nan	48
Xkinin-hobó	32
Xna-max	41
Xox	27
Xpak-um-pak	16
Xtez	15
Xtohcu	44
Xtulu	50
Xutu	39
Yaxché	30
Yaxhabin	20
Yerba mora	44
Zunumbak	44

PUBLICATIONS OF THE MUSEUM.

The series of publications of the Field Columbian Museum, of which this is a number, began with "An Historical and Descriptive Account of the Field Columbian Museum." The numbers that succeed in the series are not restricted either in authorship or subject, if coming within the scope of scientific or technical discussion, and contributive to the high standard it is hoped the publications will maintain. The series includes transactions, memoirs, monographs, bulletins, handbooks, and catalogues of collections.

The publications of the Museum are numbered consecutively, but are also sectional in character. Each series has a separate pagination and volumes. The literature of each science or general subject is in this way rendered consecutive and complete for binding.

PUBLICATIONS ISSUED:

Pub. 1. Hist. Ser., Vol. 1, No. 1. An Historical and Descriptive Account of the Field Columbian Museum.

Pub. 2. Hist. Ser., Vol. 1, No. 2. The Authentic Letters of Columbus. William Eleroy Curtis, Honorary Curator of the Columbus Memorial.

Pub. 3. Geol. Ser., Vol. 1, No. 1. Handbook and Catalogue of the Collection of Meteorites. Oliver C. Farrington, Curator of Geology.

Pub. 4. Bot. Ser., Vol. 1, No. 1. Flora of Yucatan. Charles Frederick Millspaugh, Curator of Botany.

IN PREPARATION:

Anthropol. Ser., Vol. 1, No. 1, Pt. 1. Studies among the Ancient Cities of Mexico. W. H. Holmes, Curator of Anthropology.

Anthropol. Ser., Vol. 1, No. 2, Pt. 2. Studies among the Ancient Cities of Mexico. W. H. Holmes, Curator of Anthropology.

Zool. Ser., Vol. 1, No. 1. Vertebral Column of Amia. O. P. Hay, Assistant Curator of Ichthyology.

LIBRARY
OF THE
UNIVERSITY OF CALIFORNIA